



2016

# Clinical Management of HIV In Children and Adults



## Malawi Integrated Guidelines and Standard Operating Procedures for Providing HIV Services in:

- Antenatal Care
- Maternity Care
- Under 5 Clinics
- Family Planning Clinics
- HIV Exposed Child Follow-up
- ART Clinics

**©2016 Ministry of Health, Malawi**

Publications of the Ministry of Health enjoy copyright protection in accordance with the provisions of Protocol 2 of the Universal Copyright Convention. All rights reserved.

The Ministry of Health welcomes requests for permission to reproduce or translate its publications, in part or in full.

Applications, comments and inquiries should be addressed to the Secretary for Health, P.O. Box 30377, Lilongwe 3, Malawi. We will be glad to provide the latest information on any changes made to the text, plans for new editions, and reprints and translations already available.

An electronic copy of this guideline is available on the website ([www.hiv.health.gov.mw](http://www.hiv.health.gov.mw)) of the Dept. for HIV and AIDS of the Ministry of Health.

*NOTE: The mention of certain manufacturers' products does not imply they are endorsed or recommended by the Ministry of Health in preference to others of a similar nature that are not mentioned.*



<b>Foreword to the 3<sup>rd</sup> Edition .....</b>	<b>I</b>
<b>Foreword to the 1<sup>st</sup> Edition.....</b>	<b>II</b>
<b>1 How to use these guidelines .....</b>	<b>5</b>
<b>2 Summary of new policies .....</b>	<b>6</b>
<b>3 Implementation plan .....</b>	<b>9</b>
<b>4 Integrating clinical HIV services.....</b>	<b>10</b>
<b>5 PMTCT Strategy .....</b>	<b>12</b>
<b>6 Diagnosing HIV infection and exposure.....</b>	<b>14</b>
6.1 Routine ascertainment of HIV infection status for children and adults.....	15
6.2 Routine ascertainment of HIV exposure status for children under 24 months ....	15
6.3 Presumed severe HIV disease in infants (PSHD) .....	18
<b>7 Management of HIV-related diseases .....</b>	<b>19</b>
7.1 Oral candidiasis.....	19
7.2 Oesophageal candidiasis.....	20
7.3 Chronic diarrhoea .....	20
7.4 TB .....	20
7.5 Kaposi sarcoma .....	21
7.6 Cervical cancer .....	21
7.7 Shingles (Herpes zoster) .....	21
7.8 Seborrhoeic dermatitis .....	22
7.9 Tinea corporis / cruris / pedis.....	22
7.10 Pruritic papular eruptions.....	22
7.11 Pneumocystis carinii (jiroveci) pneumonia (PCP) .....	22
7.12 Cryptococcal meningitis.....	23
7.13 Pneumonia.....	23
7.14 Sepsis .....	23
7.15 Genital ulcer disease.....	24
7.16 Urethral discharge .....	24
7.17 Abnormal vaginal discharge .....	24
7.18 Lower abdominal pain (Women, STI) .....	25
<b>8 Standard monitoring of HIV patients .....</b>	<b>27</b>
8.1 Monitoring of nutritional status .....	27
8.1.1 Children 0-14 years .....	27
8.1.2 Non-pregnant adults 15 years and above.....	27
8.1.3 Pregnant and lactating women .....	27
8.2 Standard clinical monitoring checklist.....	29

# Contents

---

8.3	CD4 count testing.....	34
8.4	Collection of DBS samples for EID and VL.....	34
<b>9</b>	<b>Preventive services for HIV patients .....</b>	<b>36</b>
9.1	Provider initiated family planning (PIFP) .....	36
9.2	Cotrimoxazole preventive therapy (CPT) .....	37
9.3	Isoniazid preventive therapy (IPT) .....	39
9.3.1	Fixed dose combination (FDC) CPT / IPT / B6.....	40
9.4	Insecticide treated bed nets (ITN).....	41
<b>10</b>	<b>Understanding ART regimens and formulations .....</b>	<b>42</b>
10.1	Classification of individual ARVs .....	43
10.2	Choosing ART regimen, formulation and dosage .....	43
10.2.1	Regimen names .....	43
10.2.2	Paediatric / adult formulations .....	44
10.2.3	Start regimen .....	44
10.2.4	Initial prescriber level .....	44
10.2.5	'Starter pack' .....	44
10.2.6	'Tail' needed .....	44
10.2.7	Contraindications.....	45
10.2.8	Adverse events / side effects.....	45
10.2.9	Dosing and frequency .....	45
10.2.10	Use of EFV in women of reproductive age .....	45
10.2.11	Use of Regimen 11 as <i>start regimen</i> for children under 3 years.....	45
10.3	Choosing regimen and time of starting in special situations.....	51
10.4	Non-standard (NS) ART regimens .....	51
<b>11</b>	<b>Prescribing and dispensing ARVs .....</b>	<b>52</b>
11.1	Rules for prescribing and dispensing of ARVs.....	52
11.2	Determining quantities to be dispensed and next appointment .....	53
11.3	Appointment / dispensing interval .....	54
<b>12</b>	<b>Starting ART .....</b>	<b>56</b>
12.1	When to start ART.....	57
12.2	Record keeping.....	57
12.3	Confirming HIV infection .....	57
12.3.1	Confirmatory testing for adults and children 2 years and above.....	58
12.3.2	Confirmatory HIV testing for children under 2 years .....	58
12.4	Preparing the patient for ART .....	61
12.4.1	Mandatory patient education when starting ART.....	61
12.5	Detecting and treating high blood pressure .....	62

12.5.1	Correct BP measurement method .....	62
12.5.2	Management of hypertension .....	63
12.6	Baseline and routine lab investigations.....	63
12.7	Combining ART and TB treatment .....	64
12.7.1	Combining 2 <sup>nd</sup> line ART with TB treatment.....	65
<b>13</b>	<b>Continuing ART .....</b>	<b>66</b>
13.1	Confirming adherence to appointment.....	66
13.2	Monitoring height and weight.....	66
13.3	Monitoring for HIV-related diseases and drug side-effects .....	66
13.4	Indications for interrupting or stopping ART.....	66
13.5	Selecting regimen and formulation for continuation.....	67
13.6	Routine TB screening ( <i>intensified case finding</i> ).....	67
13.7	Achieving optimal adherence .....	68
13.7.1	Routine adherence support .....	68
13.7.2	Intensive adherence counselling.....	69
13.8	Special treatment support for children and adolescents .....	69
13.8.1	Managing the disclosure process.....	70
13.9	Keeping track of months since ART initiation.....	71
13.10	Monitoring for treatment failure / HIV drug resistance.....	71
13.10.1	Clinical screening and diagnosis of treatment failure.....	71
13.10.2	Viral load (VL) testing .....	72
13.11	Updating follow-up outcome .....	75
13.12	Immune reconstitution inflammatory syndrome (IRIS) .....	80
13.12.1	Management of IRIS.....	80
<b>14</b>	<b>Management of labour and delivery.....</b>	<b>81</b>
14.1.1	HIV status ascertainment at maternity .....	81
14.1.2	ART provision at maternity .....	81
14.1.3	Reduce obstetric risk of HIV transmission .....	81
<b>15</b>	<b>New born care and postnatal follow-up.....</b>	<b>82</b>
15.1	Initiating integrated mother/infant follow-up .....	82
15.2	Infant and child feeding counselling.....	83
15.3	Infant NVP prophylaxis .....	84
15.3.1	Prescription and dispensing of NVP prophylaxis .....	84
15.3.2	Dosing.....	84
15.3.3	Timing and duration .....	85

# Contents

---

<b>16</b>	<b>Pre-exposure prophylaxis (PrEP) .....</b>	<b>85</b>
<b>17</b>	<b>Post exposure prophylaxis (PEP).....</b>	<b>86</b>
<b>18</b>	<b>Monitoring and evaluation.....</b>	<b>90</b>
18.1	Definitions .....	90
18.2	Reporting of registration data.....	94
18.3	Reporting of cohort outcomes .....	94
18.4	Record keeping and filing.....	95
18.5	Ensuring adequate data quality .....	96
<b>19</b>	<b>Supply Management .....</b>	<b>97</b>
19.1	HIV commodity supply cycle .....	98

## Tables

Table 1: Summary of new and updated policies .....	7
Table 2: Integrated provision and scheduling of clinical HIV services.....	11
Table 3: Schedule of HIV testing in children: Choice of type of test, interpretation of results and follow-up management.....	17
Table 4: Definition of presumed severe HIV disease (PSHD) .....	18
Table 5: Checklist for clinical monitoring of HIV exp. children and ART patients .....	29
Table 6: Detailed clinical monitoring list for HIV exp. and ART patients.....	30
Table 7: Summary protocol for preparation of DBS samples for EID and VL .....	35
Table 8: Classification of ARVs .....	43
Table 9: Standard ART Regimens (all strengths in mg).....	48
Table 10: Standard pack sizes and dosing of Paediatric and Adult formulations of ARVs, IPT and CPT.....	50
Table 11: Choosing ART regimen and timing of initiation in special situations .....	51
Table 12: Quantity of ARVs to be supplied by visit interval and daily dose .....	55
Table 13: Relevant interactions between ARVs and TB drugs.....	64
Table 14: Symptom-based identification and management of side-effects .....	76
Table 15: Dosing of NVP syrup for infant prophylaxis .....	85
Table 16: Classification of risk of transmission after exposure to HIV .....	86
Table 17: Post exposure prophylaxis regimens and dosage (number of tabs taken) .....	88
Table 18: Regimens and dose for emergency contraception.....	88
Table 19: Dosing of standard presumptive STI treatment after sexual exposure.....	89
Table 20: Overview of M&E systems for integrated HIV program reporting.....	93
Table 21: Drugs and testing supplies managed by the HIV Program .....	98

## Figures

Figure 1: Flowchart for routine ascertainment of HIV exposure / infection in children under 24 months.....	16
Figure 2: BMI classification of malnutrition for non-pregnant adults 15 years +.....	28
Figure 3: Confirmatory HIV testing for adults and children aged 2 years and above .....	59
Figure 4: Confirmatory HIV testing for children under 2 years .....	60
Figure 5: Indication, interpretation and action for routine scheduled and targeted VL testing.....	74
Figure 6: Standard follow-up schedule for HIV exposed children.....	82
Figure 7: Flowchart for HIV commodity supply management .....	99



# Acknowledgements

The Department for HIV and AIDS of the Ministry of Health gratefully acknowledges the contributions of the Technical Working Groups, under the chairmanship of Dr Frank Chimbwandira, Director of HIV Dept.:

Aziz **Abdallah**, Management Science for Health  
Allan **Ahimbisibwe**, Elizabeth Glazer Pediatric AIDS Foundation  
Saeed **Ahmed**, Baylor Pediatric Aids Initiative  
Isabel **Amoros Quiles**, Médecins Sans Frontières (France)  
Jane **Banda**, MCHIP- JHPEIGO  
Leonard **Banda**, MOH, South West Zonal Health Office  
Beth **Barr**, CDC Malawi  
Yusuf **Bhamu**, MOH, Dept. of HIV and AIDS / I-TECH  
Paul **Charlie**, Project Concern International  
Ellious **Chasukwa**, National AIDS Commission  
Thom **Chaweza**, Lighthouse Trust  
Richard **Chidakwani**, Project Concern International  
Benson **Chilima**, MOH, Public Health Reference Laboratory  
Frank **Chimbwandira**, MOH, Dept. of HIV and AIDS  
Robert **Ching'oma**, MOH, Central West Zonal Health Office  
Jacqueline **Chinkonde-Nkhoma**, UNC Project, Malawi  
John **Chipolombwe**, MOH, North Zonal Health Office  
Dorica **Chirwa**, MOH, Dept. of HIV and AIDS  
Zengani **Chirwa**, JPHIEGO  
Jonas **Chise**, MOH, Thyolo District Health Office  
Liz **Corbett**, Malawi - Liverpool Wellcome Research Programme  
Sarah **Dahoo**, Médecins Sans Frontières (Belgium)  
Luckson **Dullie**, Partners in Health  
Michael **Eliya**, MOH, Dept. of HIV and AIDS  
Daniela **Garone**, Dignitas International  
Sarah **Gibson**, PSI Malawi  
Belaineh **Girma**, MOH, NTP / I-TECH  
Dominic **Gondwe**, National AIDS Commission  
Salem **Gugsa**, Lighthouse Trust / I-TECH  
Yamikani **Gumulira**, JPHIEGO  
Sundeep **Gupta**, CDC Malawi  
Mina **Hosseinipour**, UNC Project, Malawi  
Nicolette **Jackson**, Médecins Sans Frontières (Belgium)  
Andreas **Jahn**, MOH, Dept. of HIV and AIDS  
Haswel **Jere**, DREAM Project Malawi  
Thokozani **Kalua**, MOH, Dept. of HIV and AIDS  
James **Kandulu**, MOH, Diagnostics Dept.  
Henry **Kanyerere**, MOH, National TB Control Program  
Atupele **Kapito-Tembo**, Malaria Alert Center - CDAC  
Joseph **Kasola**, MOH, Dept. of HIV and AIDS  
Blessings **Kaunda-Khangamwa**, Malaria Alert Center  
Florence **Kayambo**, JPHIEGO  
Dumbani L. **Kayira**, CDC Malawi  
Peter **Kazembe**, Baylor Pediatric Aids Initiative  
Elizabeth **Kaziya**, Peace Corps  
Maria **Kim**, Baylor Pediatric Aids Initiative  
Caroline **Kiyiika Ntale**, MOH, Dept. HIV and AIDS/ I-TECH  
Owen **Kumwenda**, USAID  
Simeon **Lijenje**, MOH, Dept. of HIV and AIDS  
Giuseppe **Liotta**, DREAM Project Malawi  
Edgar **Lungu**, UNAIDS/I-TECH  
Alice **Maida**, CDC Malawi  
Simon **Makombe**, MOH, Dept. of HIV and AIDS  
Amos **Makwaya**, MOH, Dept. of HIV and AIDS

Ken **Malisita**, Queen Elizabeth Central Hospital  
Kuzani **Mbendera**, MOH, National TB Control Program  
Takondwa **Mengezi**, Baobab Health  
Andrew **Mganga**, MOH, Dept. of HIV and AIDS  
Eustice **Mhango**, MOH, Dept. of HIV and AIDS  
Dalitso **Midiani**, MOH, Dept. of HIV and AIDS  
Chimwemwe **Mkandawire**, MOH, Dept. HIV & AIDS/ I-TECH  
Kondwani **Mkandawire**, Medical Council of Malawi  
Austin **Mnthambala**, MOH, Dept. of HIV and AIDS  
James **Mpunga**, MOH, National TB Control Program  
Nicodeme **Mubiala**, Dignitas International  
Owen **Musopole**, MOH, North Health Zonal Health Office  
Andrina **Mwansambo**, National AIDS Commission  
Charles **Mwansambo**, MOH, Chief of Health Services  
Reuben **Mwenda**, MOH, Diagnostics Dept.  
Lawrence **Nazimera**, MOH, Neno District Health Office  
Stanley **Ngoma**, MOH, Dept. of HIV and AIDS  
Joseph **Njala**, Elizabeth Glazer Pediatric AIDS Foundation  
Haldon **Njikho**, USAID  
James **Njobvuyalema**, National AIDS Commission  
Prince **Nkhoma**, Médecins Sans Frontières (France)  
Zinaumaleka **Nkhono**, Baylor Pediatric Aids Initiative  
Alice **Nkoroi**, Family Health International  
Mtemwa **Nyangulu**, CDC Malawi  
Paul **Nyasulu**, MOH, Dept. of HIV and AIDS/ I-TECH  
Goodwin **Nyirenda**, Clinton Health Access Initiative  
Mike **Nyirenda**, Partners in Hope  
Bernadette **O'Hare**, Queen Elizabeth Central Hospital  
Liesbert **Ohler**, Médecins Sans Frontières (Belgium)  
Ilhame **Ouansafi**, Clinton Health Access Initiative  
Adriana **Palomares**, Médecins Sans Frontières (France)  
Colin **Pfaff**, Partners in Hope  
Sam **Phiri**, Lighthouse Trust  
Malocho **Phoso**, Mothers 2 Mothers  
Nora **Rosenberg**, UNC Project, Malawi  
Jean Baptiste **Sagno**, DREAM Project Malawi  
Dorica **Salamba-Chirwa**, MOH, Dept. of HIV and AIDS  
Cecilia **Sambakunsi**, MOH, Dept. of HIV and AIDS/ I-TECH  
Veena **Sampamkumar**, Mothers 2 Mothers  
Abdoulaye **Sarr**, CDC Malawi  
Clara **Schlaich**, Kamuzu Central Hospital  
Alan **Schooley**, Partners in Hope  
Erik **Schouten**, Management Science for Health  
Katie **Simon**, Baylor Pediatric Aids Initiative  
Victor **Singano**, Dignitas International  
Molly **Sugme**, Partners in Health  
Dyson **Telela**, Christian Health Association of Malawi  
Rachel **Thomas**, Clinton Health Access Initiative  
Barton **Upindi**, MOH, South East Zonal Health Office  
Joep **van Oosterhout**, Dignitas International  
Anteneh **Worku**, USAID  
Emily **Wroe**, Partners in Health  
Martha **Zembeni**, Project Concern International  
Gerald **Zomba**, MOH, Dept. of HIV and AIDS/ I-TECH



## Foreword to the 3<sup>rd</sup> Edition

---

**This 3<sup>rd</sup> Edition of the *Malawi Guidelines for Clinical Management of HIV in Children and Adults* will be implemented from May 2016.** It replaces all previous editions of the Malawi Antiretroviral therapy (ART) and Prevention of Mother to Child Transmission (PMTCT) guidelines.

This document is written for medical doctors, clinical officers, medical assistants, nurses, midwives, health surveillance assistants (HSAs) and medical records clerks who are working in public and private sector health facilities in Malawi. It is designed to be a practical guide for implementation of integrated HIV Services.

The guidelines have been compiled by the joint Technical Working Groups for PMTCT, ART, HIV testing and Paediatric HIV under the leadership of the Dept. for HIV and AIDS of the Ministry of Health. The guidelines are based on ***Malawi's Revised Policy for PMTCT and ART*** which was endorsed by the Ministry of Health in June 2010 and which was prompted by the release of the 2010 Revision of the World Health Organisation (WHO) PMTCT and ART Guidelines. This 3<sup>rd</sup> Edition is an adaptation of the ***2015 WHO Consolidated Guidelines on the Use of Antiretroviral Drugs for Treating and Preventing HIV Infection***.

The protocols and policies presented in this document are adapted for health services in Malawi and follow a ***public health approach***, aiming to provide the best possible services for the largest possible number of persons in need of these services.

This document defines the framework for Malawi's National HIV Programs. Considering public health benefits and risks, as well as funding and resource implications, **deviations from these guidelines are not supported by the Ministry of Health.**

## Foreword to the 1<sup>st</sup> Edition

---

This 1<sup>st</sup> Edition of the Malawi Guidelines for Clinical Management of HIV in Children and Adults will be implemented from July 2011. It replaces all previous editions of the Malawi Antiretroviral Therapy (ART) and Prevention of Mother to Child Transmission (PMTCT) guidelines.

These guidelines are written for medical doctors, clinical officers, medical assistants, nurses, midwives, health surveillance assistants (HSAs) and medical records clerks who are working in public and private sector health facilities in Malawi. The document is designed to be a practical guide for implementation of integrated HIV services.

These guidelines have been compiled by the Joint Technical Working Groups for PMTCT, ART, HTC and Paediatric HIV under the leadership of the Department for HIV and AIDS of the Ministry of Health. The guidelines are based on Malawi's revised policy for PMTCT and ART which was endorsed by the Ministry of Health in June 2010 and which was prompted by the release of the 2010 Revision of the 2010 WHO PMTCT and ART guidelines.

The protocols and policies presented in this document are adapted for health services in Malawi and follow a public health approach, aiming to provide the best possible services for the largest possible number of persons in need of these services.

This document defines the framework for Malawi's National HIV Programs. Considering public health benefits and risks, as well as funding and resource implications, deviations from these guidelines are not supported by the Ministry of Health.

The 2<sup>nd</sup> Edition of these guidelines is scheduled for release in 2013. Any updates or amendments to protocols and policies that are to be implemented between July 2010 and the release of the 2<sup>nd</sup> Edition of the guidelines will be communicated through an official MOH circular.

## Acronyms and Abbreviations

---

<b>3TC</b>	Lamivudine
<b>AAFB</b>	Acid alcohol fast bacilli
<b>ABC</b>	Abacavir
<b>ANC</b>	Antenatal care
<b>ARM</b>	Artificial rupture of membranes
<b>ART</b>	Antiretroviral therapy
<b>ARVs</b>	Antiretroviral drugs
<b>ATV/r</b>	Atazanavir / ritonavir
<b>AZT</b>	Zidovudine
<b>B6</b>	Pyridoxine
<b>BCG</b>	Bacille Calmette-Guérin
<b>Benzyl pen</b>	Benzyl penicillin
<b>BF</b>	Breastfeeding
<b>BMI</b>	Body mass index
<b>CO</b>	Clinical Officer
<b>CPT</b>	Cotrimoxazole preventive therapy
<b>CSF</b>	Cerebrospinal fluid
<b>CTX</b>	Cotrimoxazole
<b>CXR</b>	Chest X-ray
<b>d4T</b>	Stavudine
<b>DBS</b>	Dried blood spot
<b>dl</b>	decilitre
<b>DNA-PCR</b>	Deoxyribonucleic acid polymerase chain reaction
<b>E</b>	Ethambutol
<b>EFV</b>	Efavirenz
<b>EHP</b>	Essential health package
<b>EMB</b>	Ethambutol
<b>EPI</b>	Extended Programme on Immunization
<b>EPTB</b>	Extra-pulmonary tuberculosis
<b>FDC</b>	Fixed dose combination
<b>FP</b>	Family planning
<b>GIT</b>	Gastrointestinal tract
<b>H</b>	Isoniazid
<b>Hb</b>	Haemoglobin
<b>HCC</b>	HIV Care Clinic
<b>HIV</b>	Human immunodeficiency virus
<b>HTS</b>	HIV testing services
<b>IEC</b>	Information, Education and Communication
<b>IM</b>	Intramuscular
<b>INH</b>	Isoniazid

## Acronyms and Abbreviations

---

<b>IPT</b>	Isoniazid preventive therapy
<b>IRIS</b>	Immune reconstitution inflammatory syndrome
<b>ITN</b>	Insecticide treated net
<b>IV</b>	Intravenous
<b>KS</b>	Kaposi sarcoma
<b>LFT</b>	Liver function test
<b>LPV/r</b>	Lopinavir/ ritonavir
<b>MA</b>	Medical Assistant
<b>MCH</b>	Maternal and child health
<b>MDR-TB</b>	Multi-drug resistant tuberculosis
<b>MUAC</b>	Mid-upper arm circumference
<b>NNRTI</b>	Non-nucleoside reverse transcriptase inhibitor
<b>NRTI</b>	Nucleoside reverse transcriptase inhibitor
<b>NS</b>	Non-standard (ART regimen)
<b>NVP</b>	Nevirapine
<b>OPD</b>	Out-patient Dept.
<b>ORS</b>	Oral rehydration solution
<b>PCP</b>	Pneumocystis carinii (jiroveci) pneumonia
<b>PCR</b>	Polymerase chain reaction
<b>PEP</b>	Post-exposure prophylaxis for HIV using antiretroviral medicines
<b>PI</b>	Protease inhibitor
<b>PIFP</b>	Provider initiated family planning
<b>PMTCT</b>	Prevention of mother to child transmission
<b>PO</b>	Per os
<b>PrEP</b>	Pre-exposure prophylaxis for HIV using antiretroviral medicines
<b>PSHD</b>	Presumed severe HIV disease
<b>PTB</b>	Pulmonary tuberculosis
<b>PZA</b>	Pyrazinamide
<b>R</b>	Rifampicin
<b>S</b>	Streptomycin
<b>SP</b>	Sulphadoxine / pyrimethamine
<b>STI</b>	Sexually transmitted infections
<b>TBT</b>	Anti-tuberculosis treatment
<b>TDF</b>	Tenofovir disoproxil fumarate
<b>TF</b>	Therapeutic feeding
<b>VIA</b>	Visual inspection (of the cervix) with acetic acid
<b>VL</b>	Viral load
<b>ZDV</b>	Zidovudine

# 1 How to use these guidelines

These guidelines standardise clinical management of HIV positive patients and of HIV exposed children using an integrated approach. They also incorporate relevant protocols from other national guidelines (TB, IPT, FP, STI and Reproductive Health).

Most clinical interventions for HIV patients are provided in different service delivery settings. The **standardised simplified protocols** for each intervention presented in this document will facilitate the job of the health workers and improve the standard of care for patients.



## Key Facts for Patients and Providers

- The most important information and key instructions are presented in a box at the beginning of each section.
- It is appropriate and helpful to share this information with patients during Information, Education, and Communication (IEC) sessions, and in individual counselling.

Short bullet points and 'plain language' are used throughout this document to make the information as clear and concise as possible.

## The standard package of clinical HIV interventions

**Chapter 4** on page 10 shows which of the **clinical HIV interventions** should be provided in each of the regular service delivery points of the health system. It also defines the standard package of services and explains which interventions are appropriate for which patient groups and when to deliver them.

## Protocols for how to deliver clinical HIV interventions

**Chapters 5 – 17** (page 12 – 86) explain in detail how to deliver each of the HIV interventions. The protocols and directions are the same for *all* service delivery settings. These chapters also contain several checklists, tables and flow charts which can be laminated and used as job aids in the consultation room.

## 2 Summary of new policies



### Key Facts for Patients and Providers

- All HIV infected people should start ART as soon as possible for their own health and to prevent passing the virus on to others.
- Serious HIV-related diseases can occur even in patients with high CD4 count (>500), without any previous symptoms. Immediate ART greatly reduces this risk.
- Current ART regimens are easy to take and rarely cause serious side-effects.
- ART for all HIV infected people is the most effective prevention method available: Successful ART leads to very low levels of virus in the blood and in body fluids (viral suppression). Viral suppression greatly reduces the risk of sexual or mother-to-child transmission.

There is now clear scientific evidence that patients should **start ART as soon as possible** after getting infected with HIV.

**Patient benefits:** reduced risk of serious HIV-related illnesses that can occur even in the early stages of HIV infection when the CD4 count is still above 500. Early treatment benefits outweigh the risk of short- and long-term side effects because the regimens currently used in Malawi are easy to take and usually well tolerated.

**Population benefits:** successful ART greatly reduces the risk of onward transmission to sexual partners and from mother to child.

The National Strategic Plan for HIV (2015-2020) includes the **90-90-90 treatment targets** put forward by UNAIDS, aiming to achieve viral suppression for 73% of the total HIV infected population by 2020. This will greatly reduce the number of new HIV infections and is expected to achieve epidemic control in the longer term.

The 90-90-90 strategy requires further streamlining of HIV program policies. This 3<sup>rd</sup> Edition of the Malawi Clinical HIV Guidelines aims to realize the potential for **further simplification** offered by the 2015 WHO recommendation for **universal ART eligibility**, following the same principles that were introduced in the previous editions of the National PMTCT/ART guidelines.

Table 1: Summary of new and updated policies

<i>Standard monitoring of HIV patients</i>		Page
Old	Use simplified weight for height job aid for classification of nutritional status of children.	
New	Refer to Malawi Guidelines for Community Management of Acute Malnutrition (CMAM)	27
Old	Every 3 months, do routine CD4 count for patients with confirmed HIV infection (pre-ART patients) who are not otherwise eligible for ART.	
New	Routine CD4 counts are no longer supported by the national HIV program because they do not influence routine patient management according to these guidelines.	34
<i>Preventive services for HIV patients</i>		
Old	Start isoniazid preventive therapy (IPT) at enrolment for pre-ART follow-up and continue for as long as the patient is in pre-ART follow-up. Stop IPT when ART is started.	
New	Give IPT for life to all children and adults who are receiving ART in the 10 high TB burden districts, regardless of skin test status (if known). IPT will become available as fixed dose combination tablet with cotrimoxazole and pyridoxine.	36
<i>Understanding ART regimens and formulations</i>		
Old	Use of seven standard 1 <sup>st</sup> line and three standard 2 <sup>nd</sup> line regimens. No standard 3 <sup>rd</sup> line regimen.	
New	Use of five standard 1 <sup>st</sup> line and five standard 2 <sup>nd</sup> line regimens. Stavudine-containing regimens are discontinued. Separate coding for atazanavir and lopinavir containing 2 <sup>nd</sup> line regimens. Regimen 9 is available in paediatric and adult formulation.	48
Old	Use of regimen 2P for all children started on ART.	
New	Use of protease inhibitor based regimen as 'start regimen' for children under 3 years at sites that can provide additional support. Infants under 6kg use lopinavir/ritonavir oral pellets instead of liquid formulation.	45
<i>Prescribing and dispensing ARVs</i>		
Old	Only certified clinical PMTCT/ART providers are authorized to prescribe and dispense ARVs (Doctors, Clinical officers, Medical Assistants, Registered Nurses, Nurse/Midwife Technicians)	
New	Non-health professionals (e.g. police officers) are allowed to dispense the initial dose of PEP without prior confirmation of HIV negative status if they have received PEP training by a MOH certified ART provider and if they are under regular supervision by ART clinic staff. However, continuation of PEP and ART can only be prescribed and dispensed by certified clinical ART providers.	52

**Starting ART**

Old	ART eligibility is determined by WHO clinical stage / CD4 count / age / pregnancy / breastfeeding status.	
New	Start ART as soon as possible for all HIV infected children and adults, regardless of clinical stage, CD4 count and/or pregnancy status.	<b>57</b>
Old	Children under 24 months who start ART need a confirmatory rapid antibody test at age 12 and 24 months.	
New	All children under 24 months who start ART need a confirmatory DNA-PCR. This can be collected on the day of starting ART. No follow-up testing using rapid antibody tests.	<b>58</b>
New	Screen all adults (30+ years) for hypertension at the time of ART initiation.	<b>62</b>

**Continuing ART**

New	Follow step-by-step guide for intensive adherence counselling for patients with adherence challenges and/or routine VL result >1000.	<b>69</b>
Old	Confirmation of treatment failure and switch to 2nd line requires a targeted / repeat VL result of 5000+.	
New	Targeted / repeat VL result of 1000+ AND good adherence in the 3 months before sample collection confirms treatment failure. Start 2nd line.	<b>72</b>

**Management of labour and delivery**

New	Provide a new HIV test for all women at maternity who are not already know to be HIV positive.	<b>81</b>
-----	--	-----------

**New born care and postnatal follow-up**

New	Initiate integrated mother/infant follow-up by linking the exposed child card to the mother's ART patient card.	<b>82</b>
-----	---	-----------

**Pre-exposure prophylaxis (PrEP)**

New	Other than for HIV exposed infants, the national HIV program does currently not support the use of ARVs for pre-exposure prophylaxis.	<b>85</b>
-----	---	-----------

**Post exposure prophylaxis (PEP)**

Old	Stavudine / lamivudine is used as alternative PEP regimen for children under 25kg	
New	Abacavir / lamivudine is used alternative PEP regimen for children under 25kg	<b>87</b>

**Supply Management**

Old	HIV commodities are distributed every 3 months. Facilities maintain 3-5 months of stock.	
New	HIV commodities are distributed every 2 months. Facilities maintain 2-4 months of stock.	<b>98</b>

### 3 Implementation plan

- The revised policies come into effect in **May 2016**.
- The policy changes will be communicated by circular to all District Health Offices and facilities.
- PMTCT/ART refresher trainings based on this 3<sup>rd</sup> Edition of the Clinical HIV Guidelines will commence in the first half of 2016.
- Facilities start implementing the new policies when at least 2 health workers have passed the 2016 refresher trainings.
- The next (4<sup>th</sup>) Edition of these guidelines is scheduled for release in **2018/19**. Any potential policy or protocol updates to be implemented before release of the next edition of the guidelines will be communicated through an official MOH circular.

## 4 Integrating clinical HIV services

HIV services are an integral part of the EHP. This section shows the **standard schedule** for the **minimum package of** clinical HIV interventions to be delivered within the established service points.

**Table 2** on page **11** outlines the HIV interventions to be offered at various service delivery points.

Refer to the page number for details on how to deliver the specific intervention.

### HIV Care Clinic (HCC)

- **HCC** is an integration in the same clinic setting for:
  - **HIV exposed children**
  - **ART**
- HCC services should be established in ART and MNCH clinics.
- **HCC** is designed to facilitate clinical monitoring, preventive services and ART for family members affected by HIV.
- Family appointments can be given to encourage family members to attend together for HIV services.
- Family members can be seen in the consultation room at the same time or seen individually if there are sensitive issues to discuss.

**Table 2: Integrated provision and scheduling of clinical HIV services**

Interventions that are provided only under special circumstances are marked with brackets (●)

HIV Service	Page	Schedule	OPD	In-Patients	Fam Plan Clin	ANC	Maternity	Postnatal Clin.	U5 Clinic	Exp Child FUP	ART Clinic	TB Clinic
Diagnosing HIV infection and exposure	14	Ascertain status at each visit	●	●	●	●	●	●	●	●	(●)	●
Management of HIV-related diseases	19	When diagnosed	●	●		(●)	(●)			●	●	●
Standard monitoring of HIV patients	27	At every visit								●	●	●
Provider initiated family planning (PIFP)	36	At every scheduled visit									●	
Cotrimoxazole preventive therapy (CPT)	37	At every scheduled visit				●	●			●	●	●
Isoniazid preventive therapy (IPT)	39	Dispense for 1, 2 and then 3 monthly thereafter									(●)	
Insecticide treated bed nets (ITN)	41	Dispense 1 ITN every 24 months				●			●	●	●	
Infant and child feeding counselling	83	At every visit	●			●	●		●	●	●	
Starting ART	56	As soon as possible	(●)			●	●	●			●	●
Continuing ART	66	Monthly for the 1 <sup>st</sup> 6 months; 3 monthly thereafter	(●)			●		●			●	●
Management of labour and delivery	81	On admission					●					
New born care and postnatal	82	After delivery					●	●				
Initiating integrated mother/infant follow-up	82	At first opportunity when mother known HIV+					●	●	●		●	
Infant NVP prophylaxis	84	At first opportunity when mother known HIV+				●	●	●	●	(●)		
Post exposure prophylaxis (PEP)	86	As soon as possible after risk exposure	●				●					

## 5 PMTCT Strategy



### Key Facts for Patients and Providers

- Multiple strategies are available to prevent the transmission of HIV from mother to child and to reduce the HIV burden among mothers and their children.
- These strategies are grouped into the **4 Prongs of the national PMTCT** program.
- Implemented together, these strategies have resulted in a drastic reduction of HIV infections among children. Further scale-up is expected to virtually eliminate new paediatric HIV infections and AIDS deaths among children.
- Key interventions from all 4 PMTCT prongs are covered in these guidelines, but some medical and non-biomedical interventions are beyond the scope of this document and are covered in separate guidelines.

### Prong 1: Primary prevention of HIV infection in parents

- Behaviour change communication to reduce risky sexual contacts
  - Separate strategy
- Provision of condoms
  - **Section 9.1** Provider initiated family planning (PIFP)
  - Separate condom strategy
- Voluntary medical male circumcision for HIV negative men to reduce the risk of HIV acquisition and onward transmission
  - Separate MOH guidelines
- Scale-up of HIV testing in high-yield settings for early diagnosis and ART referral
  - **Section 6.1:** Routine ascertainment of HIV infection status for children and adults
- ART provision for all HIV infected adults and children, (regardless of CD4 count and/or clinical stage) to reduce morbidity and mortality and to prevent onward transmission
  - **Section 12.1:** When to start ART
  - **Section 13.7:** Achieving optimal adherence
- Viral load monitoring and timely switch to 2nd line for patients on ART to ensure viral suppression and to reduce the risk of onward transmission
  - **Section 13.10:** Monitoring for treatment failure / HIV drug resistance
- Post-exposure prophylaxis
  - **Section 17:** Post exposure prophylaxis (PEP)

### Prong 2: Prevention of unintended pregnancies among HIV positive women

- Provider initiated family planning in ART clinics
  - **Section 9.1** Provider initiated family planning (PIFP)
  - Separate MOH guidelines: National Sexual and Reproductive Health and Rights Policy

### Prong 3: Preventing transmission of HIV from infected women to their children

- Provider initiated testing at MNCH settings for early HIV diagnosis and ART initiation
  - **Section 6.1:** Routine ascertainment of HIV infection status for children and adults
  - **Section 14.1.1:** HIV status ascertainment at maternity
- Initiation of lifelong ART for all HIV infected pregnant and breastfeeding women (regardless of CD4 count and/or clinical stage) to reduce the risk of transmission to the child.
  - **Section 12.1:** When to start ART
- Safe obstetric practices
  - **Section 14.1.3:** Reduce obstetric risk of HIV transmission
- Provision of infant nevirapine prophylaxis
  - **Section 15.3:** Infant NVP prophylaxis
- Infant feeding advice to reduce the risk transmission through breastmilk
  - **Section 15.2:** Infant and child feeding counselling

### Prong 4: Care, treatment and support for HIV-infected women and their children and families

- **Section 4:** Integrating clinical HIV services
- **Section 6.2:** Routine ascertainment of HIV exposure status for children under 24 months
- **Section 15.1:** Initiating integrated mother/infant follow-up
- **Section 7:** Management of HIV-related diseases
- **Section 9:** Preventive services for HIV patients
  - **Section 9.2:** Cotrimoxazole preventive therapy (CPT)
  - **Section 9.3:** Isoniazid preventive therapy (IPT)
  - **Section 9.4:** Insecticide treated bed nets (ITN)
- **Section 12.5:** Detecting and treating high blood pressure
- **Section 13.8:** Special treatment support for children and adolescents

## 6 Diagnosing HIV infection and exposure



### Key Facts for Patients and Providers

- **Main** HIV testing program goals:
  - **Identify as many** HIV infected people **as possible**.
  - Identify them **as early as possible** after getting infected.
  - **Ensure they start ART** as **soon** as possible.
- Additional goal is to link HIV negatives to appropriate prevention services (VMMC, etc.) and to encourage retesting based on the client risk assessment.
- **Provider Initiated Testing:** Ascertain HIV status for all patients attending health services (ANC, maternity, TB, STI, FP, U1 / U5, adult and paediatric wards).
- Remind patients during pre-test education (group or individual) that they can decline HIV testing without any 'fear of punishment' by the health worker.
- Encourage patients to attend testing with their sexual partner. Ensure that all children (including adolescents) of HIV infected parents are tested.
- Enrol all children born to and/or breastfeeding from HIV infected mothers (*'HIV exposed children'*) in the HIV Care Clinic and follow to at least age 24 months or longer if breastfeeding continues.
- From age 12 months, over 95% of children with a positive rapid test are confirmed HIV infected. Therefore, rapid testing should be used to diagnose HIV infection and start ART from age 12 months.
- Examine all children under 12 months of age with confirmed HIV antibodies for clinical conditions that constitute *Presumed Severe HIV Disease* (PSHD, see **section 6.3** on page **18**). All of these need to start ART without delay.
- All patients need a confirmatory HIV rapid test to rule out any possibility of mix-up of test results or fraudulent access to ART (also see **section 12.3** on **page 57**):
  - Before starting ART
  - Patients who received confirmatory testing at pre-ART enrolment (before 2016) do not need another confirmatory test when starting ART.
  - All children under 24 months who start ART need a confirmatory DNA-PCR using a new DBS sample. This can be collected on the day of starting ART (also see **section 12.3.2** on **page 58**).


## 6.1 Routine ascertainment of HIV infection status for children and adults

- Ask every client at every visit about the most recent HIV test and review their health passport for previous HIV test results.
- Offer HIV testing to all patients attending health facilities for any reason, if:
  - never tested
  - tested negative more than 3 months ago (follow risk assessment guidelines)
  - claims to have been tested any time in the past, but without documentation (being on ART counts as documented evidence)
- Routinely document HIV test results on page 6 of the patient's health passport unless the patient declines. For in-patients, also document test result in in-patient notes.

## 6.2 Routine ascertainment of HIV exposure status for children under 24 months

- Routinely ascertain the mother's HIV status for all children under 24 months of age seen at the U1 / U5 clinic, regardless of whether the child is healthy or sick:
  - Review mother's health passport (page 6) for the latest HIV test result
- Initiate a new HIV rapid test:
  - For the mother:
    - If she is not known to be positive and has not been tested at delivery or thereafter.
  - For the child:
    - If the mother is not available / has died
    - If the child is sick, even if the mother was tested negative during pregnancy or delivery. Mothers may have been recently infected themselves and the risk of onward transmission to the child is very high under these circumstances.
- **Figure 1** on **page 16** shows the conditions for testing of mother and/or child and the actions to be taken.

**Figure 1: Flowchart for routine ascertainment of HIV exposure / infection in children under 24 months**



**Table 3** on **page 17** shows the routine testing schedule for children under 2 years of age, the selection of the type of HIV test (DNA-PCR or rapid antibody test) depending on the child's age and the correct interpretation and action depending on the test result.

**Table 3: Schedule of HIV testing in children: Choice of type of test, interpretation of results and follow-up management**

Age (months)	Test	Schedule	Result	Interpretation	Action
<b>Under 12</b>	DNA-PCR (if available)	First opportunity from age 6 weeks	Negative	Not infected, but at risk of infection if breastfeeding	Continue HCC. Do rapid test at age 12 months.
			Positive	HIV infected	Start ART. Confirmatory DNA-PCR at ART initiation.
	Rapid antibody	Immediately if signs of PSHD identified	Negative	Not infected, but at risk of infection if breastfeeding	Treat condition. Continue HCC. Repeat rapid test at age 12 and 24 months.
		<b>OR</b> If mother's HIV status cannot be ascertained	Positive	Possibly HIV infected if no PSHD symptoms	Enrol in HCC. Do DNA-PCR at first opportunity.
				Likely AIDS if symptoms for PSHD	Start ART. Confirmatory DNA-PCR at ART initiation.
<b>12 to under 24</b>	Rapid antibody	From age 12 months	Negative	Not infected, but at risk of infection if breastfeeding	Continue HCC, repeat rapid test at age 24 m.
		<b>OR</b> If mother's HIV status cannot be ascertained	Positive	HIV Infected	Start ART. Confirmatory DNA-PCR at ART initiation.
<b>24 and above</b>	Rapid antibody	From age 24 months but ensure that BF stopped at least 6wks ago	Negative	Not infected	Discharge child from HCC.
			Positive	HIV Infected	Start ART. Confirmatory (parallel) rapid test at ART initiation.

### 6.3 Presumed severe HIV disease in infants (PSHD)

- Infants infected with HIV develop life-threatening HIV-related diseases much more quickly than older children and adults.
- It often takes too long to **confirm HIV infection** in a sick infant using **DNA-PCR**.
- Under the age of 12 months, a positive **HIV rapid antibody test does not confirm HIV infection** because maternal antibodies pass through the placenta and remain in the baby's blood for several months.
- However, a positive rapid antibody test in an infant with the following clinical signs makes **severe HIV disease** (AIDS) very likely:

**Table 4: Definition of presumed severe HIV disease (PSHD)**

Infant with positive rapid antibody test <b>PLUS:</b>		
Combination of 2:	<u>OR</u>	At least 1:
<ul style="list-style-type: none"> <li>• Oral thrush</li> <li>• Severe pneumonia</li> <li>• Severe sepsis</li> </ul>		<ul style="list-style-type: none"> <li>• Severe unexplained wasting / malnutrition not responding to treatment</li> <li>• Pneumocystis pneumonia</li> <li>• Candidiasis of oesophagus, trachea, bronchi or lungs</li> <li>• Cryptococcal meningitis</li> <li>• Toxoplasmosis of the brain (from age 1 month)</li> </ul>

- **Start ART** as quickly as possible for infants with PSHD – do not wait for a DNA-PCR result.
- Collect a DBS sample for DNA-PCR confirmatory testing on the day of starting ART (see **section 12.3** on **page 57**).

## 7 Management of HIV-related diseases



### Key Facts for Patients and Providers

- Untreated HIV infection leads to a gradual destruction of the immune system.
- Different HIV-related diseases appear at different levels of immune suppression.
- Most of these diseases can also occur in HIV negative patients, but they are a lot more common and more severe in HIV infected patients.
- Actively search and treat HIV-related diseases at ART initiation and at every follow-up visit. ART alone may not save the patient.
- Patients may have several HIV-related diseases. Write all diseases found on the ART Patient Card.
- A patient on ART who develops a new or worsening HIV-related disease may not be adherent and/or have drug-resistant HIV. Do a targeted VL after confirming good adherence in the last 3 months to rule out treatment failure. (see **section 13.10** on **page 71**)
- The risk of developing active TB is high in the first 6 months on ART and remains elevated for life (compared with HIV negative persons).
- Most HIV patients with TB do not have typical TB symptoms (productive cough) and many are sputum smear negative. It is often impossible to confirm TB diagnosis in these patients and empirical TB Treatment in very sick individuals will be lifesaving.
- HIV infected TB patients must start ART and TB treatment as soon as possible. The long term outcome is poor if only one treatment is taken.
- There is no problem with taking standard ART Regimen 5A at the same time as TB treatment.
- Use the following list to identify and manage the main HIV-related diseases seen in Malawi.

### 7.1 Oral candidiasis

#### Clinical Signs

Multiple whitish or red patches anywhere inside mouth

#### Primary Management

##### Nystatin oral suspension

Treat for 7-14 days; keep in mouth as long as possible; apply to mother's nipples if breastfeeding

**Adult:** 4ml 6-hourly

**Child:** 1ml 6-hourly

#### Secondary Management

2 Alternative treatment options if severe or no response to nystatin:

##### Fluconazole tablets

Treat for 14 days

**Adult:** 100 mg 24-hourly

**Child:** 6mg/kg on day 1 then 3mg/kg daily

##### Miconazole gum patch or gel

Use for children > 4 months and adults

Treat with 1 patch 24-hourly for 14 days

## 7.2 Oesophageal candidiasis

### Clinical signs

Retrosternal pain on swallowing; infants and children refusing to eat; +/- oral thrush

### Primary management

#### Fluconazole tablets

Treat for 14 days

**Adult:** 200mg 24-hourly for 14 days

**Child:** 12mg/kg day one then 6mg/kg

## 7.3 Chronic diarrhoea

### Clinical signs

More than 3 loose non-bloody motions per 24 hours for more than 4 weeks (adults) or 2 weeks (children)

### Diagnosis / investigations

Based on response to stepwise empirical treatment:

**Step 1** treats: isospora, cyclospora, bacterial

**Step 2** treats: giardia, clostridium, amoeba, microsporidium

**Step 3** treats: microsporidium, helminths

### Primary management

#### ORS (Thanzi)

drink 5ml/kg 4-hourly and after every episode of diarrhoea.

drink 5ml doses every 5 min if vomiting occurs

#### IV Fluids

if severe de-hydration

#### Loperamide tablets

**Adult:** 2mg after every loose stool (max 12mg in 24 hours)

**Child:** Do NOT use for children

#### Step 1: Cotrimoxazole tablets

**Adult:** 960mg 8-hourly for 7 days

**Child:** 80 mg/kg 8-hourly for 7 days

#### Zinc tablets

Give for 10 days

**Child 0-6mths:** 10 mg 24-hourly

**Child 6mths – 5 yrs:** 20 mg 24-hourly

## Secondary management

Continue with step 2 and 3 if no improvement

#### Step 2: Metronidazole tablets

**Adult:** 800mg 8-hourly for 7 days

**Child:** 15mg/kg 8-hourly for 7 days

#### Step 3: Albendazole tablets

**Adult:** 400mg 12-hourly for 14 days

## 7.4 TB

### Clinical signs

Very variable depending on organs affected.  
Persistent fever / drenching night sweats;  
weight loss; failure to thrive; cough; anaemia  
<8g/dl; enlarged nodes; meningitis signs

### Diagnosis / investigations

Often difficult to confirm in HIV+ patients.  
(Presumptive) TB case in household? 2x sputum for AAFB; Gene-Xpert; CXR; fine needle aspiration nodes (for microscopy); pleural tap for biochemistry: straw coloured effusion?  
Lumbar puncture: CSF for biochemistry, microscopy

### Primary management

#### 1<sup>st</sup> Line TB treatment

Don't delay empirical TB treatment in severely ill HIV patients with suspected TB

#### Category 1: New smear-positive or negative PTB:

Intensive phase: 2 RHZE

Continuation phase: 4 RH

#### Category 1: TB Meningitis:

Intensive phase: 2 SRHZ + prednisolone

Continuation phase: 7 RH

#### Category 2: Relapse/ return after default/ treatment failure/ recurrent TB

Intensive phase: 2 SHRZE (usually in hospital)

1 RHZE

Continuation phase: 5 RHE

### Secondary management

#### MDR-TB

Specialised treatment (see NTP guidelines)

## 7.5 Kaposi sarcoma

### Clinical signs

Single or multiple purple patches or nodes, mainly mouth, skin, conjunctiva, lung, GI tract; +/- enlarged nodes; +/- Oedema

### Diagnosis / investigations

Usually clear picture; children often present with lymphadenopathy only; consider KS even without skin or oral lesions if no response to EPTB therapy within 4 weeks

### Primary management

For all patients:

#### Analgesia

#### Symptomatic treatment

#### ART

For KS stage T0 (skin KS without oedema):

**Delayed chemotherapy** if no improvement after 3 months on ART

For KS stage T1 (KS in mouth or internal organs, nodular skin KS, skin KS with oedema):

#### Immediate chemotherapy

Contraindications for chemotherapy: Severe PN; Hb<10g/dl; platelet count <50/mm<sup>3</sup>; jaundice; pregnancy

#### 1<sup>st</sup> Line: Vincristine

Each cycle consists of 6 doses; ensure strictly IV injection as infiltration causes burns; document therapy and response in health passport; examine for recurrence at every visit

**Adult:** 2mg vincristine IV

**Child:** 0.05 mg/kg vincristine IV (max 2mg)

Review after every cycle:

Severe neuropathy / constipation: stop

Lesions cleared: stop

Good response but residual lesions: continue next cycle

Poor response: Start 2<sup>nd</sup> line chemotherapy

1) Initial cycle:

1 dose every 7 days for 6 weeks

2) Second cycle:

1 dose every 14 days for 12 weeks

3) Final cycle:

1 dose every 28 days for 6 months

### Secondary management

#### 2<sup>nd</sup> Line: vincristine + bleomycin

Cumulative max. lifetime dose for bleomycin is 400 units for adults and 17 doses for children; stop bleomycin immediately if any sign for lung fibrosis (incl. cough, shortness of breath) are seen; give one combined dose every 14 days until cumulative max. dose is reached or until response is achieved; refer for 3rd line chemotherapy (doxorubicin) if poor response

**Adult:** 15 units bleomycin IM / IV / SC **plus** 2mg vincristine IV

**Child:** 0.5 mg/kg bleomycin IM **plus** 0.05 mg/kg vincristine IV (max 2mg)

## 7.6 Cervical cancer

### Clinical signs

No early symptoms therefore active 12-monthly screening needed; abnormal vaginal discharge

### Diagnosis / investigations

#### Acetic acid visualisation (VIA)

Use good light source

Expose cervix with cusco speculum,

visualise cervix after washing for 2 minutes with a large cotton swab immersed in 4% acetic acid

### Primary management

#### Surgical, depending on stage

Cryotherapy of pre-cancerous lesions can be done immediately after VIA.

## 7.7 Shingles (Herpes zoster)

### Clinical signs

Grouped blisters in one patch; intense pain / burning; +/- fever; +/- body pains; lesions do not usually cross the body's mid-line

## Primary management

### Analgesic Ladder

Rigorous pain control

### Acyclovir tablets

Must be started before blisters burst

**Adult:** 800mg 5 times per day for 7 days

**Child:** 20 mg/kg 8-hourly for 7 days

If face affected:

Refer to Eye specialist

Monitor for secondary bacterial infection

## 7.8 Seborrhoeic dermatitis

### Clinical signs

Greasy, scaly rash in axilla, groin, scalp, neck, face

### Primary management

Clotrimazole or Miconazole cream / ointment

Hydrocortisone 1% cream/ointment

### Secondary management

#### Ketoconazole tablets

200 mg twice daily for 7 days

Flucloxacillin or Erythromycine 500mg 6-hourly for 7 days

## 7.9 Tinea corporis / cruris / pedis

### Clinical signs

Round reddened plaques with scaly edge in multiple sites, poss. widespread

### Primary management

#### Whitfield's ointment

Clotrimazole cream or Gentian-Violet paint

Apply twice daily for 3-4 weeks

### Secondary management

#### Griseofulvin tablets

**Adult:** 500 mg 12-hourly for 4-6 weeks

**Child:** 20mg/kg per day for 4-6 weeks

## 7.10 Pruritic papular eruptions

### Clinical signs

Severe itching, evenly distributed normal- or dark-coloured papules on trunk, arms or legs, often scratch-lesions

### Primary management

#### Calamine Lotion

Antihistamines

### Secondary management

#### Corticosteroid cream or tablets

#### Metronidazole tablets

250mg 12-hourly for 7-14 days

## 7.11 Pneumocystis carinii (jiroveci) pneumonia (PCP)

### Clinical signs

- Extreme shortness of breath; dry cough; +/- fever
- Severe pneumonia in infants <12 months

### Diagnosis / investigations

O<sub>2</sub> saturation: hypoxia

CXR: Diffuse interstitial or hyperinflation; bats wing shadow

Treat empirically for PCP any HIV exposed or confirmed infected infant presenting with severe pneumonia

### Primary management

#### Admit

#### Oxygen

#### Cotrimoxazole tablets

**Adult:** 4 x 480mg 8-hourly for 21 days

**Child:** 80mg/kg 8-hourly for 21 days

Lifelong maintenance (CPT)

IV Cotrimoxazole if unable to swallow and NGT impossible to place

#### Prednisolone tablets:

Give only if patient is hypoxic / in respiratory distress.

Give 15-30 minutes before cotrimoxazole

**Adult:** 8 tablets 12-hourly for 5 days  
8 tablet 24-hourly for 5 days  
4 tablets 24-hourly for 11 days

**Child:** 2mg/kg 24-hourly for 7 days  
1mg/kg 24-hourly for 7 days  
0.5mg/kg 24-hourly for 7 days

### Secondary management

#### Clindamycin

600mg 8-hourly for 3 weeks  
plus

#### Primaquine

30mg 24-hourly for 3 weeks

## 7.12 Cryptococcal meningitis

### Clinical signs

Slow onset severe headache; confusion;  
convulsions; +/- fever; +/- neck stiffness

### Diagnosis / investigations

CSF India ink stain; cryptococcal antigen in  
serum or CSF

### Primary management

#### Admit

Daily therapeutic spinal tap  
(up to 20ml per puncture)

#### Fluconazole tablets

**Adult:** 1200mg 24-hourly for 14 days  
400mg 24-hourly for 42 days  
200mg 24-hourly for life

**Child:** 12mg/kg 24-hourly for 2 weeks  
6mg/kg 24-hourly for life

### Secondary management

#### Amphotericin B

Specialised sites only

**Adult and Child:** 0.7-1mg/kg IV over 6  
hours 24-hourly for 14 days

Follow acute treatment with Fluconazole  
for life

#### Fluconazole tablets

**Adult:** 400mg 24-hourly for 42 days  
200mg 24-hourly for life

**Child:** 6mg/kg 24-hourly for life

## 7.13 Pneumonia

### Clinical signs

Productive cough; chest pain; fever; tachypnoea  
/ dyspnoea

### Diagnosis / investigations

Infiltrations on CXR

### Primary management

Child:

Mild: Tachypnoea but no dyspnoea

(See IMCI guidelines)

Adult:

Mild to moderate presentation:

#### Amoxicillin tablets

500mg 8-hourly for 5 days

**Doxycycline or Erythromycin if no response**

### Secondary management

Severe presentation:

#### Ceftriaxone 2g IV

**Add Gentamycin if no response**

## 7.14 Sepsis

### Clinical signs

Severe illness; fever (can be absent, especially in  
children); fast heart rate; fast breathing

### Diagnosis / investigations

+/- Malaria parasites; do not rule out sepsis if  
malaria parasites are seen; blood culture for  
culture and sensitivity (if available)

### Primary management

#### Health Centre Level:

Immediate presumptive treatment

Referral to hospital

Child:

**Benzyd Pen** 50,000 IU/kg IV or IM stat +

**Gentamycin** 7.5mg/kg slow IV / IM stat +

**Quinine** 10mg/kg IM stat

Adult:

**Chloramphenicol** 1g IV or IM stat +  
**Gentamycin** 240mg slow IV or IM stat +  
**Quinine** 1200mg IV in 5% dextrose over 4 hours

## Secondary management

### Hospital management:

Neonate:

**Benzyl Pen** 50,000 IU/kg IV 8-hourly +  
**Gentamycin** 7.5 mg/kg IV 24-hourly

Child:

**Gentamicin** 7.5.mg/kg 24-hourly + **Benzyl Pen** 50,000 IU/kg IV 8-hourly

OR

**Ceftriaxone** 50-100 mg/kg IV 24-hourly  
 OR (if pneumococcal sepsis suspected)

**Chloramphenicol** 25 mg/kg IV 8-hourly  
 (max. 1g per dose)

When stable continue to complete 10 days:

**Amoxicillin** 40 mg/kg (total daily dose),  
 divided into 3 doses given 8-hourly +  
**Ciprofloxacin** 15 mg/kg 12-hourly

Adult:

**Ceftriaxone** 2g IV 24-hourly

When stable continue to complete 10 days:

**Ciprofloxacin** 500 mg tablets 12-hourly +  
**Amoxicillin** 500 mg tablets 8-hourly

## 7.15 Genital ulcer disease

### Clinical signs

Skin ulcer and/or blisters on genitals with or without pain

### Diagnosis / investigations

History, examination

### Primary management

Emphasize importance of completing treatment

Avoid sex without condom until treatment complete, give min. 30 condoms

Give referral slip to treat partner

**Benzathine Penicillin**

2.4 Million Units IM stat

**Ciprofloxacin tablets**

500 mg 12-hourly for 3 days

**Acyclovir tablets**

800 mg 8-hourly for 3 days

## 7.16 Urethral discharge

### Clinical signs

Turbid discharge from urethra, usually with pain when passing urine

### Diagnosis / investigations

History, examination

### Primary management

Emphasize importance of completing treatment

Avoid sex without condom until treatment complete, give min. 30 condoms

Give referral slip to treat partner

**Gentamicin** 240 mg IM stat

**Doxycycline tabs** 100 mg 12-hourly for 7 days

**Metronidazole tabs** 2 g stat

## 7.17 Abnormal vaginal discharge

### Clinical signs

Vaginal discharge, unusual colour / odour

Itching, pain / discomfort, pain when passing urine

### Diagnosis / investigations

History, examination

### Primary management

Emphasize importance of completing treatment

Avoid sex without condom until treatment complete, give min. 30 condoms

Give referral slip to treat partner

**Gentamicin** 240 mg IM single dose

**Doxycycline tabs** 100 mg 12-hourly for 7 days

In pregnancy: **Erythromycin tabs** 500 mg 6-hourly for 7 days

**Metronidazole tabs** 1g stat

## 7.18 Lower abdominal pain (Women, STI)

### Clinical signs

Pain during sexual intercourse/ when passing urine/ around menses

Vaginal discharge / excessive bleeding at / between periods

Fever / nausea / vomiting

### Diagnosis / investigations

History, examination

### Primary management

Emphasize importance of completing treatment

Avoid sex without condom until treatment complete, give min. 30 condoms

Give referral slip to treat partner

**Gentamicin** 240 mg IM stat

**Doxycycline tabs** 100 mg 12-hourly for 14 days

**Metronidazole tabs** 400 mg 12-hourly for 14 days



## 8 Standard monitoring of HIV patients



### Key Facts for Patients and Providers

- Exposed children and ART patients need the same standard clinical assessment at every visit.
- Check actively – do not rely on patients to report problems unprompted.
- The Standard Clinical Monitoring Checklist (Table 6 on Page 30) helps to find:
  - HIV-related diseases
  - ART treatment failure
  - Drug side effects (ART, TB, CPT, IPT, etc.)
- It can be difficult to distinguish HIV-related diseases from side effects. An ambiguous symptom is likely a side-effect if it started / worsened after starting medication / improves after stopping.

### 8.1 Monitoring of nutritional status

- One of the simplest methods to detect HIV disease progression / ART treatment failure.
- Investigate any patient with weight loss for TB
- Record length / height to the nearest cm at every visit (children) / once at enrolment (adults).
- Record weight in kg to the nearest 100g at every visit (children and adults).
- Use appropriate nutrition indicator for children and adults.

#### 8.1.1 Children 0-14 years

- Classify and manage wasting / malnutrition status according to **Malawi Guidelines for Community Management of Acute Malnutrition (CMAM)**.
- Watch out for flattening of the growth curve (weight for age).


#### 8.1.2 Non-pregnant adults 15 years and above

- Classify nutrition status according to BMI from **Figure 2** on page 28
- $$BMI = \frac{weight(kg)}{height(m) \times height(m)}$$
- Watch out for any weight loss over time. Review documented previous weight whenever available as reported weight loss can be unreliable.
- BMI under 17 (below yellow curve in **Figure 2**): Start TF for 'moderate malnutrition'.
- BMI under 16 (below red curve in **Figure 2**): Start TF for 'severe malnutrition'.

#### 8.1.3 Pregnant and lactating women

- Use MUAC instead of BMI.
- MUAC less than 22cm: start TF for 'moderate malnutrition'.
- MUAC less than 19cm: start TF for 'severe malnutrition'.

Figure 2: BMI classification of malnutrition for non-pregnant adults 15 years +



## 8.2 Standard clinical monitoring checklist

- Use the summary clinical monitoring checklist to actively screen every exposed child and ART patient for clinical symptoms at every visit.
- Refer to **Table 6** on page 30 for more detailed screening instructions and interpretation of signs and symptoms for further management.

**Table 5: Checklist for clinical monitoring of HIV exp. children and ART patients**

Ask for / Examine		
Appearance: Weight loss / failure to thrive	N	Y
Body shape change / breast swelling (men)	N	Y
Swollen glands	N	Y
Headache / confusion / dizziness	N	Y
Yellow eyes	N	Y
Mouth sores	N	Y
Cough	N	Y
Shortness of breath	N	Y
Fever / night sweats	N	Y
Vomiting / abdominal pain	N	Y
Diarrhoea	N	Y
Leg pain / numbness / weakness	N	Y
Rash on arms, legs or trunk	N	Y

Table 6: Detailed clinical monitoring list for HIV exp. and ART patients

Ask / Examine	Look for	Assess	Disease (most common)	Drug Side-Effects
Appearance	<ul style="list-style-type: none"> <li>Weight loss</li> <li>Failure to thrive</li> </ul>	<ul style="list-style-type: none"> <li>Weight loss: trend from patient card / health passport</li> <li>BMI (adults)</li> <li>Weight for height, weight for age, MUAC (children)</li> </ul>	<ol style="list-style-type: none"> <li>1) TB</li> <li>2) Chronic diarrhoea</li> <li>3) Malnutrition</li> <li>4) ART treatment failure</li> <li>5) Malignancy (lymphoma)</li> </ol>	<b>Lactic acidosis due to ART</b> 1) AZT
	<ul style="list-style-type: none"> <li>Breast swelling (men)</li> <li>Body shape change</li> </ul>	<ul style="list-style-type: none"> <li>Breast enlargement (gynaecomastia)</li> <li>Slimming of cheeks</li> <li>Slimming of forearms, buttocks and legs +/- protruding veins</li> <li>Fattening of chest / belly / buttocks</li> <li>Buffalo hump</li> </ul>		<b>Gynaecomastia</b> 1) EFV <b>ART induced lipodystrophy</b> 1) AZT 2) 3TC
	Swollen glands	<ul style="list-style-type: none"> <li>Cervical / axillary lymphadenopathy</li> </ul>	<ol style="list-style-type: none"> <li>1) PGL</li> <li>2) EPTB</li> <li>3) Lymphoma</li> <li>4) KS (+/- skin lesions)</li> <li>5) BCG adenitis</li> </ol>	
Headache, confusion, dizziness	<ul style="list-style-type: none"> <li>Neck stiffness</li> <li>Nausea / vomiting</li> </ul>		<ol style="list-style-type: none"> <li>1) Meningitis (bacterial/ TB, cryptococcal)</li> <li>2) Toxoplasmosis HIV dementia</li> </ol>	<ol style="list-style-type: none"> <li>1) EFV</li> <li>2) INH</li> </ol>

Ask / Examine	Look for	Assess	Disease (most common)	Drug Side-Effects
Yellow eyes	<ul style="list-style-type: none"> <li>Yellow sclera</li> </ul>	<ul style="list-style-type: none"> <li>Jaundice</li> </ul>	1) Viral hepatitis 2) Alcoholic hepatitis 3) Malaria 4) Cancer 5) Hep B-IRIS	<b>Drug hepatitis</b> 1) NVP 2) EFV 3) PZA 4) Rifampicin 5) INH 6) Fluconazole 7) DRV/r 8) ETV <b>Hyperbilirubinaemia</b> 7) ATV/r
Mouth sores	<ul style="list-style-type: none"> <li>Mucosa lesions</li> </ul>	<ul style="list-style-type: none"> <li>Whitish patches</li> <li>Painful red patches</li> </ul>	1) Oral thrush 2) Oral hairy leukoplakia	
		<ul style="list-style-type: none"> <li>Purple lesions</li> </ul>	1) KS	
		<ul style="list-style-type: none"> <li>Ulcerations</li> </ul>	1) Acute ulcerative stomatitis/ gingivitis/ periodontitis 2) Herpes simplex 3) Angular cheilitis 4) Aphthous ulcers	<b>Hypersensitivity</b> 1) ABC 2) NVP 3) EFV 4) ETV 5) Cotrimoxazole
Cough	<ul style="list-style-type: none"> <li>Duration</li> <li>Productiveness</li> </ul>	<ul style="list-style-type: none"> <li>Less than 2 weeks</li> <li>Fever</li> <li>+/- Productive</li> </ul>	1) Pneumonia (bacterial) 2) TB suspect: circle on card 3) PCP	<b>Hypersensitivity</b> 1) ABC 2) DRV/r 3) ETV
		<ul style="list-style-type: none"> <li>More than 2 weeks</li> <li>Fever / night sweats</li> </ul>	1) TB suspect: circle on card 2) KS	

Ask / Examine	Look for	Assess	Disease (most common)	Drug Side-Effects
Shortness of breath	<ul style="list-style-type: none"> <li>Observe breathing</li> <li>Pleural effusion</li> </ul>	<ul style="list-style-type: none"> <li>Pleural effusion</li> </ul>	1) EPTB 2) Bacterial pneumonia 3) Heart failure 4) KS	
		<ul style="list-style-type: none"> <li>No pleural effusion</li> </ul>	1) Bacterial pneumonia 2) PCP 3) TB +/- pneumothorax	<b>Lactic acidosis due to ART</b> 1) AZT
	<ul style="list-style-type: none"> <li>Conjunctiva</li> </ul>	<ul style="list-style-type: none"> <li>Pale conjunctiva</li> </ul>	1) HIV anaemia 2) Chronic severe malaria 3) Nutritional anaemia	<b>Anaemia</b> 1) AZT
Fever / night sweats	<ul style="list-style-type: none"> <li>History / Duration</li> <li>Current temperature</li> </ul>	<ul style="list-style-type: none"> <li>Less than 1 month</li> </ul>	1) URTI / viral 2) Sepsis 3) Malaria 4) TB	<b>Hypersensitivity</b> 1) ABC 2) NVP 3) EFV 4) RAL 5) ETV 6) Cotrimoxazole
		<ul style="list-style-type: none"> <li>More than 1 month</li> </ul>	1) TB 2) Malignancies (lymphomas)	
Vomiting / abdominal pain	<ul style="list-style-type: none"> <li>Hydration status</li> <li>Palpate abdomen</li> </ul>	<ul style="list-style-type: none"> <li>Dehydration</li> <li>Tenderness</li> </ul>	1) TB 2) NTS sepsis 3) Acute Gastro-enteritis 4) Malaria 5) Abdominal TB 6) Ulcer disease 7) CNS disease 8) Hepatoma	<b>Drug-induced pancreatitis</b> 1) 3TC 2) RAL 3) ETV <b>Lactic acidosis due to ART</b> 1) AZT

Ask / Examine	Look for	Assess	Disease (most common)	Drug Side-Effects
Diarrhoea	<ul style="list-style-type: none"> <li>History</li> <li>Blood in stool</li> </ul>	<ul style="list-style-type: none"> <li>Less than 1 month</li> </ul>	1) Salmonella E. Coli Amoeba, Shigella 2) HIV / OI	<b>GI toxicity</b> 1) LPV/r 2) NVP 3) AZT 4) ABC 5) 3TC Antibiotics: Pseudomembranous enterocolitis
		<ul style="list-style-type: none"> <li>Longer than 1 month</li> </ul>	1) HIV / OI 2) Abdominal TB	
Leg pain, numbness, weakness	<ul style="list-style-type: none"> <li>History</li> <li>Neurological exam</li> </ul>	<ul style="list-style-type: none"> <li>Sleep disturbance (moderate)</li> <li>Motor involvement (severe)</li> </ul>	1) HIV peripheral neuropathy 2) EPTB	<b>Drug neuropathy</b> 1) INH 2) AZT 3) Vincristine 4) Metronidazole
Rash on arms, legs and trunk	<ul style="list-style-type: none"> <li>Skin lesions</li> </ul>	<ul style="list-style-type: none"> <li>Purple lesions</li> </ul>	1) KS	
		<ul style="list-style-type: none"> <li>Blisters/ vesicles</li> </ul>	1) Shingles/ varicella zoster	Stevens-Johnson Syndrome 1) NVP 2) Cotrimoxazole 3) RAL
	<ul style="list-style-type: none"> <li>Generalized rash</li> </ul>	<ul style="list-style-type: none"> <li>Maculo-papular</li> </ul>	1) HIV associated rash (PPE) 2) Fungal skin infections 3) Molluscum contagiosum 4) Scabies	<b>Skin toxicity</b> 1) NVP 2) EFV 3) CTX 4) Fluconazole 5) DRV/r 6) ETV

## 8.3 CD4 count testing



### Key Facts for Patients and Providers

- **Routine CD4 counts** are no longer supported by the national HIV program because they do not influence routine patient management according to these guidelines.
- **Targeted CD4 counts** may be requested by specialist clinicians for complicated cases.

## 8.4 Collection of DBS samples for EID and VL



### Key Facts for Patients and Providers

- Diagnosing HIV infection in infants and detecting treatment failure in patients on ART is done by testing for HIV genetic material in a blood sample.
- This requires making millions of copies of the genetic material so that there is enough to be measured. This method is called polymerase chain reaction (PCR). PCR is very sensitive and it can be disturbed by tiny amounts of dirt or contact with other samples.
- PCR testing is only done in special labs, making it necessary to prepare dried blood spot (DBS) samples that can be kept at normal temperature for several weeks.
- Carefully follow the protocol when preparing DBS samples. Most steps are the same, but there are some **important differences between** DBS for **EID** and **VL** (shown below).
- **Never** allow EID samples to **touch or mix** with VL samples as this will lead to false positive EID results:
  - Use separate rooms or at least separate tables within one room.
  - Allocate different staff for collection of DBS for EID and VL.
  - Schedule different days / time for collecting EID and VL samples.
  - Use separate drying racks, clearly labelled **EID** and **VL**.
- Pack DBS for EID and VL in separate plastic bags and envelopes.

**Table 7: Summary protocol for preparation of DBS samples for EID and VL**

	Early Infant Diagnosis (EID)	Viral Load (VL)	Caution
<b>Getting ready</b>	<ul style="list-style-type: none"> <li>Label DBS card with patient name, ID and date</li> <li>Wash hands, put on gloves, wash powder off gloves (unless powder-free)</li> </ul>		<ul style="list-style-type: none"> <li>Hold the filter paper card only at the edges</li> <li>Never touch the area near the circles</li> </ul>
<b>Sample collection</b>	<ul style="list-style-type: none"> <li>Infants &lt;9kg: select left or right side of the sole under the heel</li> <li>Children under 2 years &gt;9kg: select heel or side of big toe</li> <li>From age 2 years and adults: select side of fingertip, preferably ring finger</li> <li>Position down, warm up, squeeze intermittently</li> </ul>		
	<ul style="list-style-type: none"> <li>Wipe with alcohol swab, dry for 30 sec</li> <li>Press lancet on skin, prick, dispose into sharps bin</li> <li>Wipe away first drop of blood with sterile gauze, wait for large drop of blood to appear</li> </ul>		<ul style="list-style-type: none"> <li>Avoid excessive squeezing of heel / toe / finger</li> </ul>
	<ul style="list-style-type: none"> <li>Drip one free drop of blood directly onto filter paper</li> </ul>	<ul style="list-style-type: none"> <li>Dip capillary into blood drop and fill to black line (50 micro litres)</li> <li>Hold tip of the capillary at a slight angle in the centre of the circle on the filter paper</li> </ul>	<ul style="list-style-type: none"> <li>Don't allow the finger / toe to touch the filter paper</li> <li>Apply blood only on one side of filter paper</li> <li>Don't rub or scratch filter paper with capillary</li> </ul>
	<ul style="list-style-type: none"> <li>Let the blood soak into the paper to fill the whole circle</li> <li>Repeat this procedure until all <b>5 circles</b> are filled</li> </ul>		<ul style="list-style-type: none"> <li>Don't re-apply more blood to the same circle</li> </ul>
<b>Drying</b>	<ul style="list-style-type: none"> <li>Slot filter paper into drying rack</li> <li>Dry in protected area at room temperature for at least 3 hours (best overnight)</li> </ul>		<ul style="list-style-type: none"> <li>Don't touch/ smear/ allow to touch other objects</li> <li>Protect from sunlight, heat, dust, insects, rodents</li> </ul>
<b>Packing</b>	<ul style="list-style-type: none"> <li>Put each filter paper card into a separate zip-lock bag</li> <li>Put 3 sachets with desiccant into each zip-lock bag</li> <li>Squeeze out air and seal zip-lock bag</li> <li>Use marker pen to label the zip-lock bag and envelope, including 'EID' or 'VL'</li> <li>Insert zip-lock bags and specimen forms in this envelope and seal</li> </ul>		<ul style="list-style-type: none"> <li>Don't pack filter paper cards before completely dried</li> <li>Don't combine EID and VL samples in same envelope</li> </ul>
<b>Storage, transport</b>	<ul style="list-style-type: none"> <li>Store envelopes in cool dry place</li> </ul>		<ul style="list-style-type: none"> <li>Keep away from sunlight</li> </ul>

## 9 Preventive services for HIV patients



### Key Facts for Patients and Providers

- A simple standard package of preventive services is provided for all ART patients. This includes:
  1. Provider Initiated Family Planning (at least condoms + Depo-Provera)
  2. Cotrimoxazole Preventive Therapy
  3. Isoniazid Preventive Therapy (in districts with high TB burden)
  4. Insecticide Treated bed Nets
- This package effectively reduces:
  - HIV transmission to sexual partners
  - HIV transmission from mother to child by preventing unwanted pregnancies
- Serious HIV-related diseases (TB, diarrhoea, pneumonia, malaria, etc.)

### 9.1 Provider initiated family planning (PIFP)



### Key Facts for Patients and Providers

- Avoid unwanted pregnancies, regardless of HIV infection status.
- Use 'dual protection' – condoms alone are not enough for family planning as they have to be used very consistently.
- Sex without condom is risky if the infected partner's viral load is not suppressed. Consistent condom use is especially important in the first 6 months after starting ART and/or if viral suppression is not confirmed (e.g. low adherence and/or treatment failure).
- Some hormonal contraceptives (the pill and implants) may be less effective with ARVs or TB treatment because of drug interactions.
- Depo-Provera does not interact with ARVs or TB drugs, but it is generally slightly less effective in preventing pregnancy than implants.
- Intrauterine device, vasectomy and tubal ligation are safe to use with ART.
- Encourage HIV positive women to make an informed choice about pregnancy. Health workers should not actively discourage pregnancy as the risk of transmitting HIV to the baby is less than 5% if the mother :
  - Starts ART as early as possible, best before conception
  - Is fully adherent to ART throughout pregnancy and breastfeeding

## Implementing routine PIFP in HIV clinics

- Assume that all patients aged 15 years and above are sexually active.
- Offer condoms to all men and women age 15 years and above:
  - Minimum of **30** male and/or female condoms
- Offer counselling on contraceptive methods. Refer to FP clinic if this is not feasible in the HCC setting.
- Offer at least Depo-Provera directly in the HCC (one-stop shop)
  - 1 Depo-Provera injection every 3 months
- Give patients the opportunity to refuse either method if they feel they don't need / want it.

## 9.2 Cotrimoxazole preventive therapy (CPT)



### Key Facts for Patients and Providers

- CPT prevents PCP pneumonia, diarrhoea, malaria and other HIV-related diseases and prolongs survival.
- Start all of the following on CPT:
  - HIV exposed children from age 6 weeks
  - HIV infected children from age 6 weeks
  - HIV infected adults
- Continue CPT for life for all HIV positive patients.
- Stop CPT in HIV exposed children when confirmed negative after stopping of breastfeeding (when discharged from exposed infant follow-up).
- Provide CPT to all patients in HCC and ART follow-up.
- CPT is tolerated very well by most patients, can be taken at the same time with ART, TB treatment and IPT.
- CPT is safe in pregnancy.
- Do not combine CPT with SP – HIV positive pregnant women only take CPT (and ART).
- Children from 30.0kg and adults take one 960mg tablet of cotrimoxazole 24-hourly.
- Dispersible paediatric tablets (120mg) are used for children under 14.0kg. Dosing of paediatric CPT and ART is both based on the same weight bands.
- CPT 960mg is usually available in blister-packs of 10 tablets – 3 strips are for a 30 day supply.
- Poor adherence to CPT is a warning sign for poor adherence to ART.

## Eligibility for CPT

- All infants born to HIV infected mothers (without confirmed HIV infection) from age 6 weeks:
  - Aim to start CPT straight after the infant has finished NVP syrup.
  - Note: having taken NVP prophylaxis is NOT a condition for starting CPT.
  - Keep the infant on CPT until s/he is confirmed HIV-negative and is discharged from HCC follow-up (around age 24 months).
- Confirmed HIV infected children from age 6 weeks and adults:
  - No contra-indication against CPT in the first trimester of pregnancy.
  - Do not give SP to HIV infected pregnant women on CPT.
  - If SP has already been taken, wait for 14 days before starting CPT.

## CPT contraindications

- Jaundice
- Renal failure
- Suspected allergy to any of the following sulphonamide drugs (skin rash, mucosal ulceration, severe anaemia, leukopenia)
  - Cotrimoxazole
  - Sulfadoxine / Pyrimethamine (SP)

## CPT dosage and duration

- See **Table 10** on page **50** for dosing.
- HIV exposed children: stop CPT when confirmed HIV negative at least 6 weeks after stopping of breastfeeding.
- HIV infected children and adults continue CPT for life, unless severe side effects develop.
- Poor adherence to CPT will reduce the effectiveness of preventing HIV-related diseases, but it is less risky than poor adherence to ART.

### 9.3 Isoniazid preventive therapy (IPT)



#### Key Facts for Patients and Providers

- Daily IPT can prevent active TB disease in people who are at high risk.
- Give IPT to the following:
  - HIV infected children and adults in the 10 high burden districts (see below), regardless of TST status (if known).
  - Children under 5 years – regardless of HIV status - who live with a patient with pulmonary TB (sputum smear negative or positive; in all districts). Give 6 months course of IPT.
- Do not give IPT to a patient who has any signs suggestive of active TB: such patients need full investigation for TB and combination TB treatment if confirmed to avoid TB drug resistance.
- About 75% of all new TB cases in Malawi come from **10 high burden districts**:
  - Lilongwe, Blantyre, Mangochi, Machinga, Chikhwawa, Mzimba North, Thyolo, Mulanje, Nsanje, Chiradzulu.
  - In these districts, IPT **benefits** are expected to clearly **outweigh** the **risk** of side effects (see below).
- In the 10 high burden districts:
  - New patients: start IPT together with ART and CPT.
  - Patients already on ART: start IPT regardless of the time on ART.
  - Give IPT regardless of previous TB treatment or prior use of IPT.
  - Continue IPT for life as long as the patients remains in a high burden district.
- IPT can be taken in pregnancy and combined with CPT and ART.
- IPT is well tolerated by over 95% of patients and most side effects are mild and disappear within the first 3 months.
- Serious side effects are rare: hypersensitivity, neuropathy and severe hepatitis.
- Stop IPT if any of the following are seen:
  - Vomiting
  - Severe skin rash
  - Yellow eyes
  - Dizziness / confusion / convulsions
  - Severe numbness/burning pain and muscular weakness of legs and/or arms

## Eligibility for IPT

- Confirmed HIV infected patients in ART follow-up in one of the 10 high burden districts.
- Active TB ruled out. Use the standard TB screening questions below:
  - Current cough: any duration, productive or non-productive
  - Unexplained weight loss (adults)
  - Failure to thrive and/or malnutrition (children)
  - Fever and/or night sweat

## IPT contraindications

- Suspected or confirmed active TB
- Active hepatitis, liver damage, chronic alcohol abuse
- Severe peripheral neuropathy

## IPT dosage and duration

- See **Table 10** on page **50** for dosing.
- Give IPT during ART visits. One extra visit is needed 1 month after starting IPT.
- Review patients at month **1, 3** and **6** after starting IPT for any side-effects.
  - IPT initiation: Give INH and pyridoxine for 1 month.
  - 1 Month IPT review: Give INH and pyridoxine for 2 months.
  - From 3 Month IPT review: Continue giving INH and pyridoxine for 3 months.
- Give 1 tablet of pyridoxine 25 or 50mg 24-hourly to children and adults who are taking separate IPT (pyridoxine is already included in the FDC CPT / IPT / B6, see below).
- Continue **IPT for life** as long as the patient remains in ART care in one of the **10 high TB burden** districts.
- Stop / interrupt IPT if a patient transfers (permanently) to an ART clinic in a non-high TB burden district. Re-start IPT if the patient transfers back into a high TB burden district.
- Poor adherence to IPT will reduce the effectiveness of preventing active TB disease, but it will not cause drug-resistant TB.

### 9.3.1 Fixed dose combination (FDC) CPT / IPT / B6

- Fixed dose combination tablets of CTX 960mg / INH 300mg / B6 25mg will become available in 2016/17.
- Give the combination tablets whenever possible to reduce the pill burden and to make adherence easier.
- Patients can take 1 FDC tablet 24-hourly together with their ARVs or take ½ FDC tablet 12-hourly if they find it easier to swallow.
- Give only CTX or INH/B6 if the patient has contraindications or side effects against INH or CTX.

## 9.4 Insecticide treated bed nets (ITN)

- Dispense 1 ITN to each patient at enrolment into HIV Care.
- Dispense 1 replacement ITN every 2 years and document this on the ART patient card.

## 10 Understanding ART regimens and formulations



### Key Facts for Patients and Providers

- ART requires combining **3 different ARVs** that act differently to avoid development of drug-resistant HIV.
- Use only the standard ARV regimens for the specified patient groups shown in these guidelines. Other ARV combinations may cause more side effects or lead to drug-resistant HIV. Non-standard (NS) regimens can only be prescribed by specialists for complicated cases.
- Do not change ART regimens without clear medical indication. Unnecessary regimen changes spoil future treatment options.
- **1<sup>st</sup> Line regimens** are the best. Patients can remain on the same 1<sup>st</sup> line regimen for many years if they are fully adherent. All 1<sup>st</sup> line regimens:
  - Are easy to prescribe and easy to take.
  - Have a low risk of serious side effects and require no lab monitoring for toxicity.
  - There are **5 different 1<sup>st</sup> line** regimens:
    - **2** are standard for **initiating ART** depending on patient weight (see Table 9 on Page 48). Both are fixed-dose combinations: only 1 type of tablet is taken.
    - Move all patients with significant side effects to an alternative regimen without delay. Chose the regimen by **substituting** only the ARV responsible for the side effects.
    - All children started on Regimen 2 continue on the same regimen after their 15<sup>th</sup> birthday. Change to adult formulation (2A) above 25kg. Continue on 2A for life unless they develop toxicity or fail.
- **2<sup>nd</sup> Line regimens** are a lifeline for patients who have confirmed treatment failure on 1<sup>st</sup> line regimen (usually due to poor adherence in the past). Moving from 1<sup>st</sup> to 2<sup>nd</sup> line ART is called **switching**. 2<sup>nd</sup> line regimens:
  - Contain a completely different class of ARVs (protease inhibitors)
  - Are more complicated to prescribe and take
  - Can have more side effects
  - There are **3 different 2<sup>nd</sup> line** regimens. The appropriate 2<sup>nd</sup> line regimen is determined by the 1<sup>st</sup> line regimen that the patient was taking when failing.
  - Children **under 3 years** may respond better when **started immediately on a 2<sup>nd</sup> line** regimen. Specialized sites that can ensure extra support with giving a more complex regimen to small children should routinely initiate children under 3 years on 2<sup>nd</sup> line ART (see details on **page 45**).
- **3<sup>rd</sup> Line regimen** is a last resort for patients failing on second line in spite of good adherence. This requires confirmation of drug resistant virus using genetic analysis in the lab. 3<sup>rd</sup> line can only be initiated by a specialised ARV clinician upon authorization of review committee.
  - Very expensive
  - Can have more side effects and is more difficult to take.

## 10.1 Classification of individual ARVs

- Main classification is based on **mode of action** against HIV replication.
- Sub-classification is based on **biochemical structure** of the drug.
- Only ARVs with the same dosing interval are available as fixed-dose combinations.

**Table 8: Classification of ARVs**

Mode of action	Biochem. structure	Abbrev.	ARVs	Dosing interval
Reverse Transcriptase Inhibitors	Nucleosides	<i>NRTI</i>	AZT	12-hourly
			3TC, ABC	12- or 24-hourly
			TDF	24-hourly
	Non-Nucleosides	<i>NNRTI</i>	NVP	12-hourly
			EFV	24-hourly
			ETV	12-hourly
Protease Inhibitors		<i>PI</i>	ATV/r	24-hourly
			DRV	12-hourly
			LPV/r	12-hourly
Integrase Inhibitor		<i>II</i>	RAL	12-hourly

## 10.2 Choosing ART regimen, formulation and dosage

### 10.2.1 Regimen names

- **Table 9** shows the standard ART regimens for Malawi.
- Regimens are numbered for ease of reference:
  - Regimen 0 – 6 are 1<sup>st</sup> line regimens, including alternative 1<sup>st</sup> line regimens.
  - Regimen 1 and 3 contain stavudine (d4T) and are no longer used and have been deleted.
  - Regimen 7 – 11 are 2<sup>nd</sup> line regimens.
  - Regimen 12 is the standard 3<sup>rd</sup> line regimen.
  - An “**A**” is added to the regimen number for adult formulations (e.g. Regimen 2A) and a “**P**” is added for paediatric formulations (e.g. Regimen 2P).
- Fixed dose combinations (FDC) are shown with a slash (e.g. AZT / 3TC / NVP).
- Combinations made up of separate tablets are shown with + (e.g. AZT/3TC + EFV).
- **3TC (lamivudine)** is the backbone in **ALL** 1<sup>st</sup> and 2<sup>nd</sup> line regimens because it is extremely well tolerated and remains active even with drug-resistant HIV is present.

### 10.2.2 Paediatric / adult formulations

- Most regimens are suitable for children and adults and are available as both adult and paediatric strength tablets, but:
  - TDF may affect growing bones and is not given to children under 35kg.

### 10.2.3 Start regimen

- Start patients under 35kg on Regimen 2 and patients above 35kg on Regimen 5.
- Start on alternative 1<sup>st</sup> line regimen if the patient has any contraindications for Regimen 2 or 5.
- Start children under 3 years on regimen 11 if the site can ensure extra support with giving a more complex regimen to small children (see **section 10.2.11** on **page 45**).

### 10.2.4 Initial prescriber level

- All MOH-certified PMTCT/ART providers are authorized to start any of the seven 1<sup>st</sup> line regimens, but only experienced ART staff (certified Level 2 providers) are authorized to initiate 2<sup>nd</sup> line regimens.
- However, follow-up prescriptions for 2<sup>nd</sup> and 3<sup>rd</sup> line regimens can also be made by Level 1 providers. See details in **section 10.3** on **page 51**.

### 10.2.5 ‘Starter pack’

- Regimens with NVP (regimen **0, 2** and **6**) need to be *phased in* to avoid potentially severe hepatitis or skin toxicity. During the first 2 weeks, the NVP-containing FDC is taken only once daily (before bed). The other 2 ARVs are taken in the morning to achieve 12-hourly dosing from the first day.
- *Starter packs* are dispensed as a 2-week supply of one pack of the triple ARV fixed-dose combination (with NVP) plus one pack of the other 2 ARVs in combination (without NVP). Dispense the required number of tablets in labelled tablet dispensing bags.
- *Starter packs* are needed for all patients starting Regimen **0, 2** or **6**:
  - For the **first time** (new ART initiation)
  - After interrupting ART for more than **14 days** (re-initiation / re-start)
- *Starter packs* are **NOT** given when changing without interruption from an EFV-containing regimen (4 or 5) to regimen 0, 2 or 6. This is because patients on EFV already excrete NVP faster.

### 10.2.6 ‘Tail’ needed

- NVP and EFV remain in the body much longer than the other ARVs. Stopping any 1<sup>st</sup> line regimen due to side-effects (or due to patient’s decision, etc.) therefore requires giving a 7-day ‘tail’ of the other 2 ARVs in the regimen to avoid exposing the virus to only NVP or EFV, which would risk development of NVP- and EFV-resistant HIV and spoil future treatment options.
- However, do **NOT** give a *tail* in case of severe potentially life-threatening side effects (lactic acidosis, pancreatitis), but stop all ARVs immediately.

### 10.2.7 Contraindications

- Most contraindications are not absolute for a specific regimen: balance risks, benefits and alternatives. Usually, a suitable alternative regimen can be chosen from **Table 9**. The following conditions are absolute contraindications:
  - Patients who developed severe toxicity to any specific ARV (hepatitis or Stevens - Johnson syndrome from NVP or EFV, severe anaemia from AZT, ABC hypersensitivity) must **NEVER AGAIN** be given a regimen containing the responsible ARV.
  - Do not use **TDF**-containing regimens in severe **renal failure** (creatinine clearance <50ml/min).

### 10.2.8 Adverse events / side effects

- Chose the appropriate alternative regimen from **Alternative 1** for patients with:
  - Contraindications
  - Significant side-effects (immediately)
  - Troubling side effects that did not improve within 2 months with symptomatic treatment.
- Use **Alt. 2** if **Alt. 1** can't be used due to previous toxicity or other specific contraindications.
- The appropriate 2<sup>nd</sup> line regimen depends on the 1<sup>st</sup> line regimen the patient was on when confirmed with treatment failure. Only certified **Level 2 ART providers** can initiate 2<sup>nd</sup> line.

### 10.2.9 Dosing and frequency

- **Table 10** shows the number of tablets to be taken by children and adults once or twice per day.
- 10 weight-bands are used to determine the number of paediatric tablets to be given.
- Most paediatric formulations are **tablets** that can be crushed if necessary. The only exceptions are:
  - LPV/r and ATV/r tablets must be **given whole** (not split or crushed).
  - LPV/r for children **under 6kg** requires **liquid suspension** (80/20mg per ml) or **oral pellets** (40/10mg per capsule).

### 10.2.10 Use of EFV in women of reproductive age

- EFV is recommended in pregnancy, including in the 1<sup>st</sup> trimester. Compared with NVP, EFV provides better long-term viral suppression, has fewer adverse events and less risk of resistance.<sup>1</sup>
  - Start 5A as early as possible in pregnancy – including in the first trimester.
  - Don't change regimen if a woman got pregnant while on an EFV-containing ART regimen.

### 10.2.11 Use of Regimen 11 as *start regimen* for children under 3 years

- Children under 3 years often have a high viral load and may be infected with drug-resistant HIV from previous exposure to ARVs (mother's ART and/or infant nevirapine prophylaxis).

<sup>1</sup> Use of efavirenz during pregnancy: a public health perspective. Technical update on treatment optimization. Geneva, World Health Organization, 2012 ([www.who.int/hiv/pub/treatment2/efavirenz/en](http://www.who.int/hiv/pub/treatment2/efavirenz/en), accessed 2 December 2013).

- Therefore, Children **under 3 years** respond better when **started immediately on a 2<sup>nd</sup> line** regimen (Regimen **11**).
- Starting children on Regimen **11** requires more differentiated follow-up and mothers need more hands-on support to ensure proper swallowing and adherence to dosing:
  - Regimen **11** has a higher pill burden than the standard *Start Regimen* for children (**2**).
  - Choose the right formulation: Children **under 6kg** need LPV/r **liquid** (needs fridge, bad taste) or **oral pellets** (heat stable, taste masked). Move from LPV/r pellets to paediatric tablets as soon as the child weighs 6kg + and is able to swallow whole tabs.
  - Demonstrate how to give ARVs (see below how to give pellets) and CPT.
  - Observe regularly how the mother gives the meds. Ensure the full dose is properly swallowed.
  - Monitor VL at **6** and **12** months and every **12** months thereafter.
  - Follow-up: from age 3 years, children with confirmed viral suppression on Regimen **11** should be switched to standard 1<sup>st</sup> line (Regimen **2**). This preserves future 2<sup>nd</sup> line options and reduces the pill burden.
- Sites that can ensure the additional support (above) should routinely start all children under 3 years on Regimen **11**.
  - Don't delay ART initiation if regimen 11 is not immediately available / feasible. Start on regimen 2 instead and move to regimen 11 when possible.
- How to give **LPV/r oral pellets**:
  - Oral pellets are inside capsules. Never give the actual capsule to swallow.
  - Take out the required number of capsules and immediately close the bottle.
  - Hold the capsule on both ends and twist in opposite directions while pulling apart.
  - Empty pellets onto a clean spoon / into a feeding cup with expressed breastmilk. Immediately give to the infant.
  - Make sure the infant does not aspirate the pellets (coughing, choking, gagging).
  - Do not dissolve / crush / stir the pellets as this will release the unpleasant taste and reduce absorption.
  - Throw away the empty capsule.



**Table 9: Standard ART Regimens** (all strengths in mg)

Regimen	Paed. Formulation	Adult Formulation	Used for ART initiation 'Start regimen'	Line	Prescriber level	Starter pack	'Tail' needed	Contraindications	Possible adverse reaction	If confirmed, use	
										Alt 1	Alt 2
<b>0</b>	ABC 60 / 3TC 30 + NVP 50	ABC 600 / 3TC 300 + NVP 200	No	1 <sup>st</sup>	1	Yes	Yes	<ul style="list-style-type: none"> <li>• ABC hypersensitivity</li> <li>• Jaundice / hepatitis</li> </ul>	• Fever, body pains, vomiting, cough <sup>2</sup>	2	6, 5, NS
									• Hepatitis, rash	ABC/3TC+EFV	5, 4
									• Treatment failure	7	8
<b>2</b>	AZT 60 / 3TC 30 / NVP 50	AZT 300 / 3TC 150 / NVP 200	<ul style="list-style-type: none"> <li>• Standard for children and adults under 35kg</li> </ul>	1 <sup>st</sup>	1	Yes	Yes	<ul style="list-style-type: none"> <li>• Anaemia &lt;8g/dl</li> <li>• Jaundice / hepatitis</li> </ul>	• Anaemia, vomiting, appetite loss	0 or 5	6
									• Hepatitis, rash	4	5
									• Lipodystrophy Lactic acidosis	5	6, NS
									• Treatment failure	7	9
<b>4</b>	AZT 60 / 3TC 30 + EFV 200	AZT 300 / 3TC 150 + EFV 600	No	1 <sup>st</sup>	1	No	Yes	<ul style="list-style-type: none"> <li>• History of psychosis</li> <li>• Anaemia &lt;8g/dl</li> </ul>	• Anaemia, vomiting, appetite loss	5, 0	6
									• Lipodystrophy, lactic acidosis	5	
									• Hepatitis, rash <sup>3</sup> , psychosis, gynaecomastia <sup>4</sup>	2, 0	6
									• Treatment failure	7	9
<b>5</b>		TDF 300 / 3TC 300 / EFV 600	<ul style="list-style-type: none"> <li>• Standard for children and adults 35kg +</li> </ul>	1 <sup>st</sup>	1	No	Yes	<ul style="list-style-type: none"> <li>• History of psychosis</li> <li>• Renal failure</li> <li>• Child under 3 years <sup>5</sup></li> </ul>	• Renal failure	0 <sup>6</sup>	2 <sup>6</sup>
									• Hepatitis, rash <sup>3</sup> , psychosis, gynaecomastia <sup>4</sup>	6	0, 2
									• Persistent dizziness, visual disturbances	6	0, 2
									• Treatment failure	8	9, NS
<b>6</b>		TDF 300 / 3TC 300 + NVP 200	No	1 <sup>st</sup>	1	Yes	Yes	<ul style="list-style-type: none"> <li>• Jaundice/Hepatitis</li> <li>• Renal failure</li> <li>• Child under 3 years</li> </ul>	• Renal failure	0 <sup>6</sup>	2 <sup>6</sup>
									• Hepatitis, rash	5	NS
									• Treatment failure	8	9, NS

<sup>2</sup> Fever, body pains, vomiting, cough / sore throat and breathing problems can be due to life-threatening ABC hypersensitivity (rare). Stop all ARVs immediately. Never re-start ABC.

<sup>3</sup> Mild skin rash and/or dizziness and nightmares are common after starting EFV. This usually resolves by itself and is not usually a reason to interrupt or change regimen.

<sup>4</sup> EFV can cause breast enlargement in children and men (one side or both sides). This may resolve spontaneously while continuing EFV, but NVP substitution is usually needed (and effective).

<sup>5</sup> Regimen 5A and TDF 300mg/3TC 300mg can be used from 35kg. Paediatric formulation for TDF is currently not available in the national program.

<sup>6</sup> Patients with creatinine clearance <50 ml/min need 3TC dose reduction but full dose ABC. AZT dose needs to be reduced from CrCl <15. Combine ABC/3TC paed tabs and ABC (single) tabs to achieve the correct dose ratio. Call the HIV Dept. logistics hotline for special order of ABC (single) tabs (see page 94).

Regimen	Paed. Formulation	Adult Formulation	Used for ART initiation 'Start regimen'	Line	Prescriber level	Starter pack	'Tail' needed	Contraindications	Possible adverse reaction	If confirmed, use Alt 1	Alt 2
7		TDF 300 / 3TC 300 + ATV/r 300/100	No	2 <sup>nd</sup>	2	No	No	<ul style="list-style-type: none"> <li>Renal failure</li> <li>Patient on rifampicin <sup>7</sup></li> <li>Pre-existing jaundice or suspected hepatitis <sup>8</sup></li> </ul>	<ul style="list-style-type: none"> <li>Renal failure</li> <li>Jaundice <sup>9</sup></li> <li>Treatment failure <sup>10</sup></li> </ul>	8 <sup>6</sup>	NS
8		AZT 300 / 3TC 150 + ATV/r 300/100	No	2 <sup>nd</sup>	2	No	No	<ul style="list-style-type: none"> <li>Anaemia &lt;8g/dl</li> <li>Patient on rifampicin <sup>7</sup></li> <li>Pre-existing jaundice or suspected hepatitis <sup>8</sup></li> </ul>	<ul style="list-style-type: none"> <li>Anaemia, vomiting, appetite loss</li> <li>Lipodystrophy, Lactic acidosis</li> <li>Jaundice <sup>9</sup></li> <li>Treatment failure <sup>10</sup></li> </ul>	7	9
9	ABC 60 / 3TC 30 + LPV/r 100/25	ABC 600 / 3TC 300 + LPV/r 200/50	No	2 <sup>nd</sup>	2	No	No	<ul style="list-style-type: none"> <li>ABC hypersensitivity</li> </ul>	<ul style="list-style-type: none"> <li>Fever, body pains, vomiting, cough <sup>11</sup></li> <li>Diarrhoea, vomiting, dizziness, headache</li> <li>Treatment failure <sup>10</sup></li> </ul>	11	10
10		TDF 300 / 3TC 300 + LPV/r 200/50	No	2 <sup>nd</sup>	2	No	No	<ul style="list-style-type: none"> <li>Renal failure</li> </ul>	<ul style="list-style-type: none"> <li>Renal failure</li> <li>Diarrhoea, vomiting, dizziness, headache</li> <li>Treatment failure <sup>10</sup></li> </ul>	11 <sup>6</sup>	8 <sup>6</sup>
11	AZT 60 / 3TC 30 + LPV/r 100/25	AZT 300 / 3TC 150 + LPV/r 200/50	<ul style="list-style-type: none"> <li>Preferred start regimen for children under 3 years at sites with extra support (page 45)</li> </ul>	2 <sup>nd</sup>	2	No	No	<ul style="list-style-type: none"> <li>Anaemia &lt;8g/dl</li> </ul>	<ul style="list-style-type: none"> <li>Anaemia, vomiting, appetite loss</li> <li>Lipodystrophy, lactic acidosis</li> <li>Diarrhoea, vomiting, dizziness, headache</li> <li>Treatment failure <sup>10</sup></li> </ul>	10	7
12		DRV 600 + r 100 + ETV 100 + RAL 400	No	3 <sup>rd</sup>	2	No	No	<ul style="list-style-type: none"> <li></li> </ul>	<ul style="list-style-type: none"> <li>Diarrhoea, vomiting, headache, dizziness</li> <li>Neuropathy</li> <li>Rash</li> <li>Jaundice</li> </ul>	NS	NS
<div> <div>3TC</div> <div>Lamivudine</div> </div> <div> <div>ABC</div> <div>Abacavir</div> </div> <div> <div>ATV</div> <div>Atazanavir</div> </div> <div> <div>AZT</div> <div>Zidovudine</div> </div> <div> <div>DRV</div> <div>Darunavir</div> </div> <div> <div>EFV</div> <div>Efavirenz</div> </div> <div> <div>ETV</div> <div>Etravirine</div> </div> <div> <div>LPV</div> <div>Lopinavir</div> </div> <div> <div>NVP</div> <div>Nevirapine</div> </div> <div> <div>r</div> <div>Ritonavir</div> </div> <div> <div>RAL</div> <div>Raltegravir</div> </div> <div> <div>TDF</div> <div>Tenofovir</div> </div>											

<sup>7</sup> Do not combine ATV/r with rifampicin (TB treatment). Substitute to LPV/r for the duration of TB treatment.

<sup>8</sup> Do not start patients with pre-existing jaundice or suspected hepatitis on ATV/r. Use LPV/r instead.

<sup>9</sup> ATV/r can cause jaundice. Mostly, this is only of cosmetic concern. Refer jaundice to a specialist for LFT. If only indirect bilirubin is raised, continue ATV. Stop ATV/r if LFT cannot be done.

<sup>10</sup> Treatment failure on 2<sup>nd</sup> line ART needs confirmation of resistance mutations by genotyping before 3<sup>rd</sup> line ART can be considered.

<sup>11</sup> Fever, body pains, vomiting, cough / sore throat and breathing problems can be due to life-threatening ABC hypersensitivity (rare). Stop all ARVs immediately. Never re-start ABC.

**Table 10: Standard pack sizes and dosing of Paediatric and Adult formulations of ARVs, IPT and CPT**

Drug	Tablets per tin		3 – 3.9 kg		4 – 5.9 kg		6 – 9.9 kg		10 – 13.9 kg		14 – 19.9kg		20 – 24.9kg		25 – 29.9kg		30 – 34.9 kg		35 – 39.9 kg		40 kg +	
	Paed.	Adult	AM	PM	AM	PM	AM	PM	AM	PM	AM	PM	AM	PM	AM	PM	AM	PM	AM	PM	AM	PM
NVP	60	60	1	1	1	1	1½	1½	2	2	2½	2½	3	3	1	1	1	1	1	1	1	1
AZT / 3TC	60	60	1	1	1	1	1½	1½	2	2	2½	2½	3	3	1	1	1	1	1	1	1	1
AZT / 3TC / NVP	60	60	1	1	1	1	1½	1½	2	2	2½	2½	3	3	1	1	1	1	1	1	1	1
ABC / 3TC	60	60	1	1	1	1	1½	1½	2	2	2½	2½	3	3	0	1	0	1	0	1	0	1
LPV / r liquid / tabs	60	120	1ml	1ml	1.5ml	1.5ml	2	1	2	1	2	2	2	2	3	3	3	3	2	2	2	2
LPV / r pellets (in caps)	120		2	2	2	2																
EFV	90	30							0	1	0	1½	0	1½	0	2	0	2	0	1	0	1
ATV / r		30																0	1	0	1	
TDF / 3TC		30																0	1	0	1	
TDF / 3TC / EFV		30																0	1	0	1	
DRV		60																1	1	1	1	
r		60																1	1	1	1	
ETV		120																2	2	2	2	
RAL		60																1	1	1	1	
CTX 120	1000		0	1	0	1	1	1	1	1	2	2	2	2								
INH 100	100		0	½	0	½	0	1	0	1½	0	2	0	2½								
CTX 480		1000					0	½	0	½	0	1	0	1	0	2	0	2	0	2	0	2
CTX 960		1000									0	½	0	½	0	1	0	1	0	1	0	1
INH 300		672									0	½	0	½	0	1	0	1	0	1	0	1
CTX960/ INH300/ B6 25		30									0	½	0	½	0	1	0	1	0	1	0	1

### 10.3 Choosing regimen and time of starting in special situations

Table 11: Choosing ART regimen and timing of initiation in special situations

Condition	Timing of ART initiation	Weight	
		Less than 35kg	35kg +
<b>Anaemia (&lt;8g/dl)</b>	<ul style="list-style-type: none"> <li>As soon as possible</li> </ul>	<b>0P</b>	<b>5A</b>
<b>Active TB</b>	<ul style="list-style-type: none"> <li>Within 14 days of diagnosis</li> <li>TBT + ART can be started on the same day if the patient is stable. Don't delay TBT or ART</li> </ul>	<b>Under 3 yrs: 2P</b> <b>3 yrs +: 4P / 4A</b>	<b>5A</b>
<b>Jaundice</b>	<ul style="list-style-type: none"> <li>Refer to District or Central Hospital</li> <li>After investigation and stabilisation</li> </ul>	<b>4P / 4A</b>	<b>5A</b>
<b>1<sup>st</sup> trimester pregnancy</b>	<ul style="list-style-type: none"> <li>As soon as possible</li> </ul>		<b>5A</b>
<b>In labour (new HIV+)</b>	<ul style="list-style-type: none"> <li>As soon as possible</li> </ul>		<b>5A</b>
<b>Renal failure</b>	<ul style="list-style-type: none"> <li>Refer to District or Central Hospital</li> <li>Start within 7 days of diagnosis</li> </ul>	<b>0P / 0A</b>	<b>0P / 0A</b>
<b>Psychiatric Illness (history)</b>	<ul style="list-style-type: none"> <li>As soon as possible</li> <li>Reliable guardian needed</li> </ul>	<b>2P / 2A</b>	<b>6A</b>

### 10.4 Non-standard (NS) ART regimens

- Only expert ART clinicians can initiate NS regimens.
- Patients with multiple contraindications and/or adverse reactions against all standard NRTIs (TDF, AZT, ABC) or NNRTIs (NVP, EFV) may need a NS regimen.
- Consider ATV/r or LPV/r for substitution of NVP and EFV.
- Contact the Dept. for HIV and AIDS for availability of non-standard ARVs (see **section 19** on **page 97**).
  - Provide patient history, indication and proposed regimen.

# 11 Prescribing and dispensing ARVs



## Key Facts for Patients and Providers

- ARVs should be taken after the same number of hours every day (e.g. every 12 or every 24 hours). Most ART regimens can be taken in the morning, at noon or at night and it does not matter if they are taken before, after or with food.
  - EFV (regimen 4 and 5) can cause dizziness, especially in the first 4 weeks. This is less troublesome when taken before bed.
- **Missing a dose:** what to do if a patient remembers to take his ARVs late? If the patient remembers:
  - **Less than half-way** to the next scheduled dose: take the missed dose immediately, and take the regular next dose at the normal time.
  - **More than half way** to the next scheduled dose: skip the missed dose and take the regular next dose at the normal time.
- Dispense ARVs only in the original sealed container. Only exception: open containers to dispense the precise number of tablets needed for *Starter Packs*.<sup>1</sup>
- Only the patient or his registered guardians/treatment supporter is allowed to collect ARVs.
- In an emergency, patients are allowed to collect ARVs from any ART clinic in Malawi following special rules (see below).

## 11.1 Rules for prescribing and dispensing of ARVs

### ARVs for treatment of HIV (ART)

- Only MOH-certified clinical ART providers are authorized to **prescribe** ART: Medical Doctors; Clinical officers; Medical Assistants; Registered Nurses; Nurse/Midwife Technicians.
- Only health workers and qualified pharmacy personnel are allowed to **dispense** ARVs.
- ARVs may not be further distributed outside of certified ART health facilities.
- Only the patient or his individual registered guardian/treatment supporter is allowed to **collect** ARVs.

### ARVs for PEP

- PEP needs to be started as soon as possible after high risk exposure. Such events are often managed under challenging circumstances (e.g. rape, accidents).
- Non-health professionals (e.g. police officers) are allowed to dispense the initial dose of PEP without prior confirmation of HIV negative status under the following circumstances:
  - Received PEP training by a MOH certified ART provider.
  - Under regular supervision by ART clinic staff

- DHOs are responsible for supplying and accounting for ARVs given to e.g. *Victim Support Units* and must provide active support with ARV stock management.

### Emergency dispensing to patients from another PMTCT/ART site

- In an emergency, patients are allowed to collect ARVs from any ART clinic in Malawi under the following conditions:
  - The patient must present an ART identity card or the health passport with ARV dispensing information.
  - If in doubt about a patient's authenticity, confirm by calling the site where the patient is registered.
  - Document emergency ARV dispensing in the patient's health passport.
  - ARV dispensed to patients registered at another site must be recorded in the *Emergency ARV Dispensing Register*. Improve a hardcover register: Date, original ARV registration number, original facility name, patient name and contact details, ARV name and quantity dispensed, reason for emergency dispensation, staff name.
  - Instruct patient to return to their ART clinic of registration as soon as possible to ensure the patient is not recorded as defaulter.

## 11.2 Determining quantities to be dispensed and next appointment

- **Table 12** on page 55 shows the number of tablets to be supplied for appointment intervals of 2, 4, 8 or 12 weeks for the total number of tablets taken of each ARV per day (paediatric and adult formulations).
  - Use **Table 10** to add up the 'total tablets taken per day' for each ARV contained in the regimen. For example: a child of 15kg on AZT/3TC/NVP (Regimen 2) takes **2½** paediatric tablets in the morning and **2½** tablets in the evening, adding up to **5** total tablets per day.
  - The *Actual number of tablets needed* is the minimum number of total tablets the patient needs to take home to cover the time to the next appointment. (Total tablets = tablets remaining from the previous visit + tablets newly dispensed). The number needed includes an extra 2-day supply to act as a safety-buffer. The total tablets must meet or exceed the *Actual number of tablets needed*.
  - Different ARVs come in tins of 30, 60, 90 or 120 tablets (see **Table 10**). Given that only full tins should be dispensed, the number of tablets needed is *rounded up* to multiples of full tins.
  - *Rounding up* may result in a considerable *over-supply*. For some regimens and dosages, perfectly adherent patients will be left with more than half a tin of ARVs at their next appointment. Explain this to the patient / guardian and emphasize the importance of keeping the next appointment.
  - The number of tablets expected to be used in the interval is shown for 'perfect adherence' (100%) and for 'good adherence' (95%-105%).
  - Calculate the number of tablets used by subtracting total tablets remaining at the current visit from total tablets available at the end of the previous visit.

### 11.3 Appointment / dispensing interval

- Give next appointment date at least 2 days before ARVs would be finished to allow for the safety buffer.
- Take account of the weekly ART clinic schedule (e.g. Mondays + Wednesdays) when giving the next appointment. Appointments are usually given for 2 weeks (starter pack), 4, 8 or 12 weeks.
- Patients initiating standard or alternative first line ART have to be reviewed clinically after **2 weeks** if they have been given a starter pack / otherwise after **1 month** and then every month for the first 6 months.
- Thereafter, stable and adherent patients can be given up to 12-week (**3-month**) appointments.
- In exceptional cases (e.g. international travel), up to **6** or even **12 months** of ARVs can be dispensed.
- Patients starting 2<sup>nd</sup> line ART have to be seen every 4 weeks for the first 6 months. Thereafter, patients who are stable and adherent to 2<sup>nd</sup> line ART can be given up to 8-week appointments.
- Align dispensing of CPT and IPT with ART visits.
- Push back appointment date to allow patients to use up accumulated 'hanging' tablets, e.g. give an appointment after 5 instead of 4 weeks.

Table 12: Quantity of ARVs to be supplied by visit interval and daily dose

<b>Note:</b> supply and consumption must be calculated <u>separately for each component</u> in the regimen. Example: separate calculation for AZT/3TC and AZT/3TC/NVP making up a starter pack of Regimen 2												
Dispens. interval	Total tabs taken per day	Supply needed									Total tabs <b>USED</b> in interval - Adherence	
		Actual tabs *	<i>Multiples of full tins</i>									
			Tins of 30		Tins of 60		Tins of 90		Tins of 120		Perfect	Good
			tabs	tins	tabs	tins	tabs	tins	tabs	tins	100%	95% – 105%
2 weeks	1	16	30	1	60	1	90	1			14	14 – 14
	1 ½	24			60	1	90	1			21	20 – 22
	2	32			60	1	90	1			28	27 – 29
	2 ½	40			60	1					35	34 – 36
	3	48			60	1					42	40 – 44
	4	64			120	2					56	54 – 58
	5	80			120	2					70	67 – 73
	6	96			120	2					84	80 – 88
4 weeks	1	30	30	1	60	1	90	1			28	27 – 29
	1 ½	45					90	1			42	40 – 44
	2	60			60	1	90	1			56	54 – 58
	3	90			120	2			120	1	84	80 – 88
	4	120			120	2			120	1	112	107 – 117
	5	150			180	3					140	133 – 147
	6	180			180	3			240	2	168	160 – 176
	8	240			240	4					224	213 – 235
	9	270			300	5					252	240 – 264
8 weeks	1	58	60	2	60	1	90	1			56	54 – 58
	1 ½	87					90	1			84	80 – 88
	2	116			120	2	180	2			112	107 – 117
	3	174			180	3			240	2	168	160 – 176
	4	232			240	4			240	2	224	213 – 235
	5	290			300	5					280	266 – 294
	6	348			360	6			360	3	336	320 – 352
	8	464			480	8					448	426 – 470
	9	522			540	9					504	479 – 529
12 weeks	1	86	90	3	120	2	90	1			84	80 – 88
	1 ½	129					180	2			126	120 – 132
	2	172			180	3	180	2			168	160 – 176
	3	258			300	5					252	240 – 264
	4	344			360	6					336	320 – 352
	5	430			480	8					420	399 – 441
	6	516			540	9					504	479 – 529

\* Actual tabs needed includes a 2-day safety-buffer

## 12 Starting ART



### Key Facts for Patients and Providers

- ART does not cure HIV infection.
- ART stops the virus from multiplying, which allows the immune system to recover.
- The virus will 'wake up' as soon as ART is interrupted and it will learn how to evade ART. This means that ART may no longer work for this patient.
- Once started, ART must be taken every day for life. All patients need effective support:
  - Identify a reliable guardian / treatment supporter who needs to attend ART education.
  - Link with patient / peer support group
- Successful ART leads to very low levels of virus in blood, semen and vaginal fluids. This greatly reduces the risk of sexual or mother-to-child transmission. However, condom use is important
  - In the first 6 months after starting ART
  - Later if adherence is not good and/or viral suppression has not been confirmed.
- All patients need a confirmatory HIV test before starting ART (see **section 12.3** on **page 57**) to rule out any possibility of mix-up of test results or fraudulent access to ART.
- ARVs must not be dispensed outside of certified PMTCT/ART facilities (static or outreach) and must not be shared, sold or passed on to others.
  - Bring back any remaining ARVs at every clinic visit to allow the provider to count them.
  - Return unused ARVs (e.g. after a patient's death) to the clinic for proper disposal.
- Patients who are late for their ART appointment will be actively followed from the clinic (home visit, phone, guardian).
  - Ask for consent for active follow-up at the time of starting ART.
  - Patients can withdraw consent at any time.
- A small number of patients on ART develop serious side-effects. Educate all patients about the important signs to look out for (see *Key facts* on **page 61**)

## 12.1 When to start ART



### Key Facts for Patients and Providers

- Start ART as soon as possible:
  - For all children and adults with **confirmed HIV infection**
  - For infants with **presumed AIDS** (following definition of PSHD).
- **All** patients need a confirmatory HIV test before starting ART.
- Explain the benefits of immediate ART for the patient's **own health**, and for **prevention** of onward transmission to sexual partners and from mother to child. This understanding is key for patient motivation and good adherence.
- Patients may not be ready to start ART immediately.
  - Allow for reflection time if the patient is unsure and/or wants to discuss with family.
  - Schedule a follow-up appointment not further than 2 weeks.

## 12.2 Record keeping

- PMTCT/ART nurse or clinician: fill ART patient cards immediately when ART eligibility is established (do not delegate this to HSA). For this reason, keep blank ART treatment cards at OPD, ANC, maternity, wards, etc.
- Dispensing of ARVs must be recorded on the patient treatment cards.
- Complete ART treatment cards before giving out the first supply of ARVs.
- Patients should only be entered in the ART register when receiving their first supply of ARVs.

## 12.3 Confirming HIV infection

- All patients need a confirmatory HIV antibody test to rule out a mix-up of test results or fraudulent access to ART:
  - Before starting ART
  - Patients who received confirmatory testing at pre-ART (before 2016) do not need another confirmatory test when starting ART.
  - All children under 24 months who start ART need a confirmatory DNA-PCR using a new DBS sample. This can be collected on the day of starting ART
- Do not delay ART initiation if HIV test kits are not available for the confirmatory test, but do confirmatory test at the next scheduled visit as soon as testing is available.

### 12.3.1 Confirmatory testing for adults and children 2 years and above

- Attach a dedicated HIV testing provider to the ART clinic to do confirmatory testing. Ensure that all Quality Assurance protocols for HIV testing (proficiency testing, quality control) are being followed.
- **Use the first and second rapid test in parallel** (currently Determine + Uni-Gold) for confirmatory HIV testing. Review **Figure 3** on **page 59** for the correct algorithm and interpretation of results:
- Test 1 and Test 2 are **both positive**:
  - Record '**Confirmatory Positive**' result in the **MOH HIV Rapid Testing register** (Version 3, January 2013)
  - Record confirmatory HIV test results on ART patient card.
  - Start ART
- Test 1 and Test2 are **discordant**:
  - Review testing protocol, quality control, expiry date and condition of test kits. Do immediate (parallel) repeat. Use a different (experienced) HIV testing provider if possible.
    - Both positive: see above
    - Discordant or both negative: see below
- Test 1 and Test2 are **both negative**:
  - Record '**Confirmatory Inconclusive**' in HIV Testing Register.
  - Collect DBS blood sample and send to reference lab and/or send patient to referral hospital to repeat regular HIV testing and for review by an experienced ART clinician.
  - Give follow-up appointment to review lab test result.

### 12.3.2 Confirmatory HIV testing for children under 2 years

- All children to be started on ART under the age of 2 years need a confirmatory DNA-PCR.
- Collect the DBS sample on / before the day of initiation.
- Don't delay ART initiation - don't wait for the confirmatory PCR result before starting ART.
- Review **Figure 4** on **page 60** for the schedule of follow-up testing and the correct action based on the results.

Figure 3: Confirmatory HIV testing for adults and children aged 2 years and above





Figure 4: Confirmatory HIV testing for children under 2 years



\* Give NRTI 'tail' when interrupting ART regimen OP or 2P (see **section 10.2.6 on page 44**).

## 12.4 Preparing the patient for ART

- Start ART as soon as possible after testing positive.
- Offer pregnant women to start ART on the same day of diagnosis.
- Confirm that patient (or parent/guardian if patient is <15 years) understands implications of ART and is committed to lifelong adherence.
- Identify long-term treatment support for patients who are unable to take responsibility for their own treatment (persons with mental disability or drug-addiction, etc.).
- Ask all patients to attend the initial group counselling and/or the ART initiation visit with a named guardian/treatment supporter.
  - Another patient can be appointed as the *named treatment supporter* if the patient is unable to identify a suitable guardian.

### 12.4.1 Mandatory patient education when starting ART



#### Key Facts for Patients and Providers

- A small number of patients on ART develop significant side-effects.
- Most side-effects are mild and disappear while ART is continued.
  - EFV can cause **bad dreams** and **dizziness** in the first few weeks of treatment, but this usually disappears by itself and it is important to continue treatment.
- Some side-effects require a regimen change.
  - Ask all men/boys to monitor themselves for **swelling of the breast** (gynaecomastia). Report this at the next scheduled visit. Substitute EFV with NVP as soon as possible after onset of gynaecomastia to improve likelihood of full reversal.
- Very few patients develop serious side effects. Stop all drugs immediately and present to the hospital if any of the following conditions are seen:
  - **Yellow eyes / hepatitis**
  - Severe **stomach pain** and **vomiting**
  - Severe **skin rash** with blisters, involving eyes, mouth or genitals

- All patients must receive individual counselling at ART initiation.
- Women starting ART in labour can receive individual ART counselling after delivery.
- In addition, all patients should attend an ART group counselling session. Recommended practice:
  - Attended group counselling between 1 to 5 days before the day of ART initiation.
  - But: group counselling can be on the same day as initiation to avoid delay beyond 7 days.
  - Pregnant women may attend the group counselling at the next scheduled visit to ensure they can start ART on the same day.
  - Ask patients to attend with their named guardian (also see **section 12.3** on **page 57**).

## ART group counselling

- Use the latest version of the MOH ART flip chart.
- Share “Key facts for providers and patients”
- Explain the standard VL monitoring schedule (see **page 72**). Ask the patients to help remember when VL is due.

## Individual ART counselling

- Confirm that patient and guardian have understood the following:
  - Commitment to lifelong adherence
  - Dosage and interval of taking ARVs
  - Potential side-effects
  - Date of next appointment

## 12.5 Detecting and treating high blood pressure



### Key Facts for Patients and Providers

- 1 out of 3 adults in Malawi have hypertension and over 90% of these have not been diagnosed.
- Even without hypertension, HIV patients have a higher risk of stroke.
- Treating all hypertensive ART patients can prevent many cases of stroke, heart and kidney failure and other complications.
- Screen all adults (**30 years +**) for hypertension:
  - At least once at the time of ART initiation. Record BP on patient card header.
  - Aim to repeat BP screening at least every 12 months

### 12.5.1 Correct BP measurement method

- Make sure the patient is relaxed (rest at least 5 minutes after physical activity).
- Sit upright, remove clothing from upper arm that may restrict blood flow or interfere with BP cuff.
- Make sure BP cuff is the right size: check the arm circumference is within range shown on the cuff.
- If the initial reading is higher than **140** systolic and/or **90** diastolic:
  - Repeat reading twice. Wait for at least 5 minutes between readings.
  - Calculate the average between the 3 readings (separately for the systolic and diastolic values).

Classification	Systolic		Diastolic	Management
<b>Mild</b>	140-159	<i>and/or</i>	90-99	Try <i>lifestyle measures</i> alone, start stepped treatment if no normalization
<b>Moderate</b>	160-179	<i>and/or</i>	100-109	<i>Lifestyle measures</i> + stepped treatment
<b>Severe</b>	>180	<i>and/or</i>	>110	Urgent treatment <i>Lifestyle measures</i> + stepped treatment

### 12.5.2 Management of hypertension

- Start management for hypertension if the average of the 3 readings is higher than **140** systolic and/or **90** diastolic.
- Urgent treatment for severe hypertension if repeat reading is **180** systolic and/or **110** diastolic.
- Fill NCD patient card for monitoring and documentation.
- Screen for diabetes
- *Lifestyle measures*: Eat more veg and fruit, less meat / fat, reduce salt, stop smoking, exercise regularly, normalize weight, limit alcohol
- See **Malawi Standard Treatment Guidelines 2015** for stepped anti-hypertension treatment.

## 12.6 Baseline and routine lab investigations

- The national program does not require:
  - Routine baseline lab investigations before starting ART or routine investigations for ART toxicity.
  - Routine scheduled CD4 monitoring of patients on ART is not supported.
- Use targeted investigations if clinically indicated.
- Scheduled VL monitoring is being rolled out (see **section 13.10** on page **72**).

## 12.7 Combining ART and TB treatment



### Key Facts for Patients and Providers

- Each year, 19,000 (2%) of the 1,000,000 HIV infected Malawians develop active TB and 7,000 die from TB<sup>12</sup>.
- The risk of developing active TB is particularly high for the first 6 months on ART and remains elevated for life.
- Most HIV patients with TB do not have typical TB symptoms (productive cough) and many are sputum smear negative.
- HIV infected TB patients must start ART and TB treatment as soon as possible. The long term outcome is poor if only one treatment is taken.
- There is no problem with taking ART Regimen 5A at the same time as TB treatment.

- Certain combinations of ARVs and TB drugs increase the risk of side-effects or reduce each other's effectiveness (due to faster excretion).
- The following table shows the relevant interactions.
  - **Green:** Combination causes no problems
  - **Yellow:** Combination causes usually no problems but monitor patient for possibly increased side-effects or adjust dosage as shown
  - **Red:** Do not combine without specialist advice

**Table 13: Relevant interactions between ARVs and TB drugs**

	Isoniazid	Rifampicin	Streptomycin	Ethambutol	Pyrazinamide
<b>TDF</b>	OK	OK	renal toxicity	OK	OK
<b>AZT</b>	OK	OK	OK	OK	OK
<b>3TC</b>	OK	OK	OK	OK	OK
<b>EFV</b>	OK	OK	skin rash	OK	hepatitis
<b>NVP</b>	skin rash	start NVP full dose, hepatitis	skin rash	OK	hepatitis
<b>ABC</b>	OK	OK	OK	OK	OK
<b>ATV/r</b>	OK	no experience (don't combine)	OK	OK	OK
<b>LPV/r</b>	OK	major dose adjustment	OK	OK	OK

<sup>12</sup> 2015 Global tuberculosis report (WHO)

### 12.7.1 Combining 2<sup>nd</sup> line ART with TB treatment

- Patients with ART failure (see **section 6.9.9** on **page 60**) may develop active TB. In this case, second line ART needs to be combined with TB treatment.
- Do not combine ATV/r with rifampicin-containing TB treatment. Give LPV/r instead of ATV/r for the duration of TB treatment and move (back) to ATV/r once TB treatment has been completed.
- Double the daily dose of LPV/r (4 tablets of LPV 200mg / r 50mg every 12 hours) for the duration of rifampicin treatment.
- Alternatively, replace rifampicin with rifabutin in patients on LPV/r (normal dose). Give rifabutin 150mg daily. Other TB drugs in the regimen should also be continued.

## 13 Continuing ART

### 13.1 Confirming adherence to appointment

- On the patient card, look at the *Next Appointment Date* given at the previous visit to confirm that the patient is not late.
- The patient is likely to have missed doses if s/he is more than 2 days late. Compare and validate with *Pill Count* and the reported number of *Doses Missed*.

### 13.2 Monitoring height and weight

- Record current weight (and height for children under 18 years).
- Look for weight changes compared with previous measurements. Patients are expected to normalize their weight in the first 6-12 months on ART.
- Classify nutrition status for children based on *CMAM* guidelines.
- Investigate any consistent weight loss over 2 or more consecutive visits. Remember to confirm that the scale is correctly calibrated and any heavy clothing was removed.

### 13.3 Monitoring for HIV-related diseases and drug side-effects

- Use the standard clinical monitoring checklist for HIV patients to actively screen for symptoms of HIV-related diseases and/or drug side effects.
- Use the syndromic guide shown in **Table 14** on **page 76** to identify the likely cause of symptoms and to choose the right primary and secondary management.
- A symptom that could be caused by an HIV-related disease or by a side-effect is more likely a side-effect if it started or worsened after the start of medication.
- Circle side-effects Yes / No on the patient card and specify new side effects under *Notes*.
- Change the ART regimen if medically indicated (see below).
- Write any new HIV-related disease under *Notes* on the back of the patient card.

### 13.4 Indications for interrupting or stopping ART

- Stop ART in patients with chronic poor adherence. Consider stopping if 3 intensive counselling sessions have failed.
- ART should be stopped abruptly and completely if any of the following severe side-effects are suspected:
  - Lactic acidosis
  - Pancreatitis
  - Severe hepatitis
  - Stevens-Johnson syndrome

- Stopping ART in patients with less severe toxicity against EFV or NVP (skin rash, psychiatric effects) should be done by giving a 'tail' of the other 2 ARVs for 7 days to prevent 'monotherapy' due to the long half-life of NVP and EFV (see **Table 9** on **page 48**).

## 13.5 Selecting regimen and formulation for continuation

- Don't change regimen without clear medical indication. Unnecessary changes spoil future treatment options.

### Do NOT change ART regimen:

- If a patient has moderate dizziness / drowsiness / nightmares in the first 2-4 weeks of starting a regimen with EFV (regimen 4 or 5. Also see footnote on **page 48**).
- All children on Regimen 2 continue on the same regimen after their 15<sup>th</sup> birthday. These patients continue on Regimen 2A through adolescence and adulthood unless they develop toxicity or fail.

### Change dosage and formulation:

- Review current weight for children and adjust dosing if necessary. Children on 1<sup>st</sup> line regimens change to adult formulation and dosage when their weight is over 25kg (see **Table 10** on **page 48**).
- Start a new ART Patient Card – Adult ARV Formulations for children who change from paediatric to adult ARV formulation. File together with the old card in the same polythene sleeve.

### Change ART regimen:

- Use **Table 9** on **page 48** to select the appropriate alternative regimen. Change patients with significant side-effects immediately. Change patients with troubling side-effects that did not improve after 2 months of symptomatic treatment.
- Children who were on paediatric 2<sup>nd</sup> line regimen (Regimen 9P) routinely change to standard adult 2<sup>nd</sup> line regimen (Regimen 7A) once they weigh over 35kg. This is to reduce the pill burden while continuing on an equally effective regimen.
- Add any new regimen to the *ART Regimens* history section on the card header and specify any non-standard regimen here.
- Patients with multiple contraindications and/or side-effects may need a NS regimen (see Section 10.4 on **page 51**)

## 13.6 Routine TB screening (*intensified case finding*)

- Screen all patients at each visit for signs of active TB using 4 standard screening questions
  - Cough of any duration
  - Fever
  - Night sweats
  - Weight loss / failure to thrive / malnutrition

- Classify screening outcome as follows:
  - **TB not suspected** if none of the 4 signs are positive. In this case, the patient is very unlikely to have active TB.
  - **TB suspected** if one or several of the 4 signs are positive.
    - Thoroughly investigate further (full clinical exam, sputum for AAFB, chest x-ray, fine needle aspirate, etc.).
    - Interrupt IPT until active TB has been ruled out
  - **TB confirmed** if the patient has a current confirmed episode of TB (clinical or lab diagnosis).
    - Always confirm if the patient is currently taking and adherent to TB treatment – initiate TB treatment without delay or provide intensive adherence support.
    - Classify **on TB treatment** or **not on treatment**.

## 13.7 Achieving optimal adherence



### Key Facts for Patients and Providers

- Patients must take more than 95% of doses at the prescribed interval for life to prevent HIV drug-resistance. Repeated skipping of individual doses or repeated longer interruptions inevitably lead to development of HIV drug-resistance.
- **Example:** HIV drug-resistance will develop if a patient on Regimen 5A (TDF/3TC/EFV) continues to skip more than **3 tablets** in every **8 week** period.
- Children and adolescents on ART need special support (see **page 69**).

### 13.7.1 Routine adherence support

- Ask at every clinical assessment visit:
  - Have you had any problems taking your ARVs? Can you explain what problems?
  - Were there any days when you did not manage to take all of your tablets at the right time? (Weekends, weekdays, mornings, evenings?)
- Remind patients of the importance of perfect adherence at every clinic visit:
  - Initial ART counselling
  - Follow-up group counselling
  - Start *intensive adherence counselling (IAC)* if any sign for poor adherence (see **page 69**)
- Give **practical strategies** how to achieve optimal adherence:
  - Build ARVs into the daily routine (e.g. before washing the face, after evening meal)
  - Ask family or friends to remind
  - Set a daily alarm on the cell phone

- Keep a 'drug diary' and mark every tablet taken
- Encourage honest dialogue. Avoid giving the impression of 'policing' the patient. Work with patients to help them achieve good adherence.
- Poor adherence always has valid reasons and most can be resolved: vomiting, transport problems, domestic problems, (perceived) side effects, psychological problems, wrong understanding, etc.

### 13.7.2 Intensive adherence counselling

#### Indications

- Questionable or confirmed poor adherence noted at regular visit / late for appointment
- Routine VL result > 1,000 (suspected treatment failure)

#### Step-by-step guide

- Ask both patient and the treatment supporter to attend.
- Explain the information presented in the boxes with **Key facts**:
  - Starting ART (page 56)
  - Achieving optimal adherence (page 68)
  - Monitoring for treatment failure / HIV drug resistance (page 71)
- Make a (verbal) **contract** with the patient and the treatment supporter:
  - "We will check your VL again in 3 months."
  - "We will work together to make sure that the result will be **VL undetectable**."
- Identify the specific problems / situations that get in the way of good adherence. Ask for:
  - Frequent travel
  - Conflicts at home
  - Alcohol / drug problems
  - Mood disorder / depression
- Agree on an action plan and write instructions in health passport: select the most suitable **practical strategies** from **page 68**. Review specific strategies for children / adolescents (**page 69**)
- Give monthly appointments until follow-up VL is due (after 3 months of good adherence).
  - Do pill count and assess adherence closely at each follow-up visit
  - Review action plan: what has worked – what has not? Revise plan if necessary.

### 13.8 Special treatment support for children and adolescents

- Ask at every visit:
  - Who is responsible for supervising the taking of ARVs?
  - Who stands in for the guardian if s/he is away?
  - How do you give the tablets?

- Select a teacher or fellow student as treatment supporter for children attending boarding school.
  - Offer to transfer the child to the most convenient ART site closest to school.
- Children (just like any other patients) who are adherent and stable on ART can be given 3 months of drug supply or more if necessary.

### 13.8.1 Managing the disclosure process

- Explain to the parent that disclosure is a gradual process. Assure the parent that you will work through this process together.
- Remind parents / care givers at every clinic visit that it is very important to talk to the child about their HIV infection and ART status.
- Don't isolate the child behind a "wall of secrecy and silence". Remember the child probably knows more than you think.
- Never lie or make up stories about the child's HIV infection and the drugs they are taking (e.g. misrepresenting ARVs as TB drugs or vitamins). Lies will eventually come out and undermine trust and make the child feel guilt, shame and will damage self-esteem and lead to poor adherence.
- Ask parents at every visit how far they have come in the disclosure process.
- Encourage parents to talk directly to their child in the environment they feel most comfortable. Offer to take part in the discussion if parents are uncomfortable doing this on their own.

#### From age 5-7 years:

- Explain that the child has a germ that requires taking drugs every day to keep the germ 'asleep'.

#### By age 11-13 years:

- Add more information gradually. By age 11-13 years, all of the following should have been explained:
  - Touching, cuddling and kissing are safe.
  - Sharing soap, towel, plates and cutlery is safe.
  - Don't share needles or razor blades. HIV and other diseases can travel in traces of blood and infect the other person.

#### From puberty / adolescence:

- Invite open dialogue about 'teenage challenges' that can get in the way of good adherence:
  - Low self esteem
  - Conflicts at home / at school
  - Relationships
  - Alcohol / drug abuse
- Link up with *Teen Club* or other peer support groups
- Offer condoms; explain use on penis model; give at least 20 condoms

- Explain: Don't have penetrative sex without condom. HIV can travel in semen and vaginal fluid and infect the other person.
- Explain: It is still possible for you to have children later in life. The risk of passing HIV to your partner or to your baby is very low if you are taking your ARVs every day as prescribed.
- Explain: Where to access STI treatment, family planning services and help in case of sexual assault.

## 13.9 Keeping track of months since ART initiation

- Needed to determine when blood samples for routine VL monitoring are to be drawn.
- Calculate and document on the ART patient card the number of months since the patient first started ART. Simply calculate the number of months since first ART initiation, ignoring any potential gaps (periods of stopping / defaulting).
- Electronic medical record systems give automatic reminders when scheduled VL samples are due.

## 13.10 Monitoring for treatment failure / HIV drug resistance



### Key Facts for Patients and Providers

- ARV drug resistance starts gradually and the virus will still be partly suppressed for many months. Emerging drug-resistant virus does not cause any immediate clinical symptoms.
- HIV will grow resistant to more and more ARVs if a patient continues to take a failing ART regimen for several months. Accumulated multiple ARV resistance can make it difficult to find a second line regimen that still works.
- HIV drug resistance usually affects different ARVs of the same class.
- **Example:** HIV that has grown resistant to EFV will also be resistant to NVP, even if the patient has never taken NVP before.
- Drug resistant virus can be transmitted to other people.
- **Example:** About 5% of Malawians who got newly infected with HIV in 2009 acquired virus with some level of drug-resistance against d4T/3TC/ NVP.

### 13.10.1 Clinical screening and diagnosis of treatment failure

- **Suspect ART failure** if both of the following clinical conditions are met:
  - On ART for 12 months or more
  - New HIV-related disease / unexplained weight loss / failure to thrive
- For all **suspected ART failure** cases, look for indications for poor adherence in the last 6 months
  - Adherence was good:
    - Do a targeted VL or refer to have this done immediately.
  - Adherence was questionable:
    - Start intensive adherence counselling (see **page 69**)

- Do a targeted VL after 3 months if adherence was satisfactory.
- See **Figure 5** on **page 74** for the interpretation of VL results.

### 13.10.2 Viral load (VL) testing



#### Key Facts for Patients and Providers

- VL is the best measure for the level of progression of HIV infection.
  - VL = number of viral particles per ml of blood.
  - More virus ⇒ faster destruction of CD4 cells ⇒ more severe immunosuppression.
- Successful ART leads to such low levels of HIV in the blood that it can no longer be detected with VL testing. An **undetectable VL** is also called **viral suppression**. This is the **aim of ART**.
- VL testing is expensive.
- VL testing uses an advanced lab method (RNA-PCR) on a blood sample. It can be done from:
  - Dried blood spot (DBS): Transport in plastic bag with desiccant at ambient temperature, sample viable for 3 months or more (see **section 8.4** on **page 34**).
  - Blood plasma: Transport in cooler box to lab within 24 hours.
- VL is required to confirm suspected ART failure (clinical and/or CD4-based).
- Routine VL monitoring is being scaled up gradually.
- The VL schedule is designed to detect ART failure early while avoiding unnecessary tests to save cost:
  - Patients with drug-resistant HIV when starting ART may have a high VL after **6 months** on ART. This can be from infection with drug-resistant HIV or after taking sdNVP. Otherwise, a high VL at 6 months is an important sign for poor adherence.
  - After that, patients who are adherent and well have a low risk of ART failure. Therefore, routine follow-up VLs are done at **2 years, 4 years, 6 years, etc.** after ART initiation.
- Do additional targeted VLs outside of this schedule when suspecting ART failure.
- Explain the standard VL monitoring schedule to every patient. Ask the patient to help remember when VL is due.
- Actively communicate (phone / home visit) VL results >1000 to patients as soon as the result is received at the site. Call for an early appointment.

#### When to do VL

- **Routine scheduled** VL is done for all patients at specific times after ART initiation:
  - At 6 months, 2 years, 4 years, and every 2 years thereafter.
  - Collect catch-up VL sample at the next opportunity if the regular schedule was missed. Continue with the regular schedule (determined by the time since ART initiation).

- **Targeted/Repeat**

- Routine VL result was 1,000+ and patient has received IAC for 3 months and we are confident adherence in the last 3 months was good.
- Patient with clinically suspected treatment failure and we are confident adherence in the last 3 months was good.
- Mandatory before starting 2<sup>nd</sup> line ART to confirm suspected ART failure.

## Interpreting and acting on VL results

- Review **Figure 5** on **page 74** for indication, interpretation and action from VL testing.

### Successful ART

<b>Finding</b>	<b>Routine or targeted / repeat VL below detection limit or less than 1,000</b>
<b>Interpretation</b>	Successful ART
<b>Action</b>	Praise the patient and encourage further good adherence. Continue on the same regimen. Monitor VL at next milestone.


### Potential treatment failure

<b>Finding</b>	<b>Routine VL result 1,000+</b>
<b>Interpretation</b>	Potential treatment failure
<b>Action</b>	Start intensive adherence counselling (see page 69). Continue on the same regimen. Collect repeat VL after 3 months of good adherence.

### Confirmed treatment failure

<b>Finding</b>	<b>Targeted / repeat VL result 1,000+ <u>AND</u> good adherence</b> in the 3 months before sample collection
<b>Interpretation</b>	The virus is likely resistant to the current ART regimen.
<b>Action</b>	Start / continue intensive adherence counselling. <u>Patients on 1<sup>st</sup> line:</u> consult certified 2 <sup>nd</sup> Line Prescriber for initiation of 2 <sup>nd</sup> line ART without delay. <u>Patients on 2<sup>nd</sup> line:</u> consult certified 2 <sup>nd</sup> Line Prescriber to consider genotype resistance testing. Call the HIV Dept. hotline (see page 102) to organize resistance testing. Request 3 <sup>rd</sup> line ARVs if resistance to 2nd line ART has been confirmed. 'Reset the clock' for routine VL monitoring: 6, 24 months, etc. after switch to 2nd or 3rd line.

Figure 5: Indication, interpretation and action for routine scheduled and targeted VL testing



\* Any of the following: Significant unintended weight loss, failure to thrive, new or worsening HIV-related disease (suspected or confirmed)

### 13.11 Updating follow-up outcome

- Regularly review all patient cards and keep an appointment register to identify patients who are overdue for their appointment as soon as possible.
- Try to contact the patient or the named guardian by phone or by home visit from 2 weeks after the missed appointment. Confirm from ART Patient Card that consent was given for home visit.
  - Patient is alive: counsel to return to the clinic as soon as possible and continue treatment.
  - Patient has stopped, died or transferred out: update outcome and date of outcome on patient card and in register.
- Loss to follow-up ('default'):
  - Patient is overdue for the appointment and is not known to have stopped ART, died or transferred to another facility.
  - Classify as 'defaulted' if the patient has run out of ARVs 2 or more months ago (based on the number of tins given at the last visit).
- Patients who are alive but known to have stopped ART (for any reason) should be classified as 'stopped' and not as 'defaulted'.
- Ask guardians to notify the clinic if an ART patient has died. Bring back the patient health passport and/or ART ID and any remaining ARVs.

Table 14: Symptom-based identification and management of side-effects

Cause (in order of likelihood)	Diagnosis	Primary Management	Secondary Management
<b>Body pains, weakness</b>			
AZT, 3TC	Severe anaemia: Hb <7 g/dl	Stop AZT, consider transfusion	Substitute AZT, continue ART without gap
AZT	Lactic acidosis (LA): shortness of breath, nausea Serum lactate: suspect: 2-5 mmol/l, confirmed: ≥5 mmol/l	<i>Any suspected LA:</i> Stop all ART immediately IV fluids, treat at hospital	Don't re-start ART before lactic acid <2mmol/l Can re-start ART with AZT after <u>suspected</u> LA Never give AZT after <u>confirmed</u> LA Can use ABC or TDF containing regimen
<b>Fever</b>			
Onset independent of drugs: Bacteraemia, malaria	FBC, MPs, blood culture, urine dipstick		
Onset within 8 weeks of starting drugs: ABC, NVP, EFV	ABC, NVP or EFV hypersensitivity: Body pains, vomiting, diarrhoea, abdominal pain, sore throat, cough, shortness of breath, rash, jaundice	Any suspected hypersensitivity: Stop all ART immediately, treat at hospital	Do not re-start before symptoms have resolved Never use NVP or ABC again Replace NVP with EFV and ABC with TDF
<b>Slimming: Cheeks, forearms, buttocks, legs (often prominent veins) Fattening: Back of neck ('buffalo hump'), breast, stomach, and waist</b>			
AZT, LPV/r, 3TC, TDF, HIV EFV	Lipodystrophy (from ART / HIV itself)	Reassure patient Substitute likely causative ARV	
<b>Breast swelling / enlargement: one- or both-sided, in males or children</b>			
EFV, ketoconazole, cimetidine, omeprazole, spironolactone, isoniazid testosterone deficiency (HIV), AZT, LPV/r	Gynaecomastia: palpate enlarged breast gland Lipodystrophy: accumulation of fat (from ART / HIV itself)	Reassure patient Substitute EFV with NVP in ART regimen.	Consider surgery for extreme gynaecomastia

Cause (in order of likelihood)	Diagnosis	Primary Management	Secondary Management
<b>Upper GI symptoms: Nausea, vomiting</b>			
AZT, LPV/r, 3TC	Lactic acidosis ? (see ' <i>Body pains and weakness</i> ') Jaundice? (see ' <i>Yellow eyes</i> ')	Adults only: Promethazine 25 mg up to 12-hourly. Adults or children (lower dose): Chlorpheniramine (Piriton) 10 mg up to 8-hourly-oral rehydration solution(ORS)	If no lactic acidosis: try to continuing the same ART regimen If persistent, substitute
<b>Skin Rash</b>			
Onset before starting drugs: Seborrhoeic dermatitis ("bumpy itch")	HIV-related skin rash	Adults only: Promethazine 25 mg 12-hourly Adults or children (lower dose): Chlorpheniramine (Piriton) 10 mg 8-hourly Calamine lotion	Consider scabies, etc.
Onset within 8 weeks of starting drugs: NVP, ABC, Cotrimoxazole, EFV	Mild hypersensitivity Macular/popular rash <u>not</u> involving mouth, eyes, and genitalia No fever, body pain, weakness, etc.	Continue EFV, reassure: initial rash mostly resolves. Continue on half dose NVP (if on NVP starter pack) for further 2 weeks Adults only: Promethazine 25mg 12-hourly Adults or children (lower dose): Chlorpheniramine (Piriton) 10 mg 8-hourly	If no improvement on half dose NVP, stop NVP Substitute to EFV once rash has resolved. If patient unable to take EFV, consult with ART specialist for alternatives
<b>Lower GI symptoms: Diarrhoea, lower abdominal pain</b>			
Onset before ART initiation: HIV-induced	Stepwise empirical treatment	Stepwise empirical treatment of chronic HIV diarrhoea (see page 20)	
Onset within 6 weeks of starting drug: LPV/r, AZT, 3TC	Drug toxicity	For adults only: Loperamide 2 mg 8-hourly (mainly for LPV/r induced diarrhoea)	Try to continue same ART regimen If persistent substitute

Cause (in order of likelihood)	Diagnosis	Primary Management	Secondary Management
<b>Severe upper abdominal pain, nausea and vomiting</b>			
3TC	Pancreatitis Serum amylase >1.5 times above upper normal limit	Stop all ART immediately Treat at hospital	Restart ART after complete remission Use TDF- or AZT-containing regimen
NVP, EFV, alcohol, viral hepatitis	Acute fulminant liver failure Liver function tests	Discontinue ART immediately Treat at hospital Identify cause and manage accordingly	Never re-start NVP or EFV if this was the suspected cause Reinitiate ART one month after jaundice is resolved, and LFT <2.5 of upper normal limit
<b>Yellow eyes</b>			
Viral hepatitis, alcohol, ATV/r, NVP, INH, EFV, ABC, severe malaria, cancer	LFT and ultrasound scan to differentiate: Viral hepatitis, cirrhosis, drug hepatitis, primary liver cancer, metastases	Discontinue ART immediately if jaundice develops on ART. See footnote 9 on page 49 for patients on ATV/r. Identify cause and manage accordingly (LFT, ultrasound, hepatitis serology).	Never re-start NVP or EFV if this was the suspected cause. Re-initiate ART 1 month after jaundice has resolved and LFT <2.5 times upper normal limit.
<b>Swollen face and eyelids, particularly in the morning/tiredness, too much or too little urine</b>			
Onset before starting drugs HIV, diabetes, hypertension	Confirm nephropathy with serum creatinine	Identify cause and manage accordingly	Adjust ART dosage according to creatinine clearance
Onset within 1 year of starting drugs: TDF, streptomycin	Confirm nephropathy with serum creatinine	Admit to hospital Substitute TDF to AZT without gap Stop streptomycin	Adjust ART dosage according to creatinine clearance

Cause (in order of likelihood)	Diagnosis	Primary Management	Secondary Management
<b>Drowsiness, confusion, nightmares, psychosis</b>			
EFV	Neuropsychiatric EFV toxicity	Drowsiness/ bad dreams usually disappear after a few weeks without the need to discontinue ART. Take EFV before bed. Confusion / psychosis: replace EFV with NVP immediately	If intolerable replace EFV with NVP
<b>Leg pain, numbness or burning, inability to walk</b>			
Onset before starting drugs: HIV neuropathy	Mild peripheral neuropathy (PN): no sleep disturbance	Amitriptyline 25 mg nocte for 4 weeks Pain control using WHO analgesic ladder	If no improvement after 4 weeks: stop amitriptyline, continue analgesics
Onset or worsening after starting drugs INH, vincristine	Moderate PN: sleep disturbance	Stop responsible drug WHO analgesic ladder	
Onset independent of drugs Alcohol, diabetes	Severe PN: severe pain, muscular weakness		

## 13.12 Immune reconstitution inflammatory syndrome (IRIS)



### Key Facts for Patients and Providers

- A small number of patients may get worse in the first 6 months after starting ART.
- The most common causes for this are (in the order of likelihood):
  - Undiagnosed / untreated OI, mainly TB
  - Poor adherence to ART
  - Drug-resistant TB (if on TB treatment)
  - IRIS
- IRIS is an over-aggressive response of the immune system caused by a sudden recovery on ART.
- IRIS appears as a severe bout / worsening of an OI:
  - TB
  - Cryptococcal meningitis
  - Herpes zoster
  - KS
  - Hepatitis
- IRIS should only be considered if the more common causes for worsening have been ruled out.
- Patients who start ART with very advanced AIDS are at a higher risk of developing IRIS.
  - Recent / concurrent treatment for TB or cryptococcal meningitis.

### 13.12.1 Management of IRIS

- Confirm that ART is actually taken as prescribed.
- Continue ART if ART toxicity has been ruled out as the underlying cause.
- Treat the OI.
- Consider TB treatment failure if worsening occurs after more than one month on TB treatment.
- Admit severe cases to hospital.
- Seek specialist advice on whether NSAIDs and/or prednisolone should be given.

## 14 Management of labour and delivery

### 14.1.1 HIV status ascertainment at maternity

- Review HIV testing page in health passport on admission.
- Provide **new HIV test**<sup>13</sup> for all women, who are:
  - Not already known to be HIV positive
  - Never tested or tested negative any time in the past, even if this result is from the last trimester.

### 14.1.2 ART provision at maternity

- Mothers already on ART: continue the same ART regimen at regular prescribed intervals. Pregnancy / breastfeeding are no indication to change women from any previous ART regimen.
- HIV positive mothers not yet on ART / who interrupted / stopped ART: emergency ART initiation
  - Start lifelong TDF/3TC/EFV (Regimen 5A) as soon as possible, during labour or after delivery.
  - Deliver individual ART counselling and IEC before discharge.

### 14.1.3 Reduce obstetric risk of HIV transmission


- Use a partogram to allow early detection and management of prolonged labour.
- Artificial rupture of membranes (ARM) increases the risk of HIV transmission.
  - ARM is not indicated if labour is progressing well.
  - If prolonged labour due to poor uterine contraction: perform ARM at ≥6cm cervical dilation and augment with oxytocin (pitocin).
- Do not perform routine episiotomy except for specific obstetric indications (e.g. vacuum extraction).
- Avoid frequent vaginal examinations.
- Do not 'milk' the umbilical cord before cutting.
- Do not suction with a naso-gastric tube unless there is meconium-stained liquor.
- Immediately after birth, wipe the baby dry with a towel to remove maternal body fluids.

<sup>13</sup> There is no general time limit for offering HIV testing. Consider that other important interventions such as C-section or tubal ligation are also offered with emergency counselling very late in labour.

## 15 New born care and postnatal follow-up

- Follow regular post-natal care.
- Give BCG and oral polio vaccine after birth to all babies born to HIV infected mothers (as for all other infants).
- See **Figure 6** below for the standard schedule of HIV exposed child follow-up: NVP prophylaxis, CPT, feeding and HIV testing

**Figure 6: Standard follow-up schedule for HIV exposed children**



### 15.1 Initiating integrated mother/infant follow-up

- Ensure continued follow-up for HIV infected mothers and babies.
- Enrol baby in HCC before discharge from post-natal ward:
  - Fill Exposed Child patient card, enter in HCC register.
- Mothers on ART before delivery:
  - Confirm next ART appointment.
  - Synchronise mother's ART appointment with baby's first HCC visit. Aim for HCC enrolment at post-natal visit or first vaccination visit.
- Mother initiated ART in labour:
  - Fill ART patient card and enter in ART register.
  - Write baby's HCC registration number on mother's ART card.
  - Give regular 4 week ART + HCC appointment.
- If mother wants to continue HCC and ART at another facility:
  - Record 'transfer out' in HIV clinic and ART register and give mother her ART patient card and the baby's Exposed child card.

## 15.2 Infant and child feeding counselling



### Key Facts for Patients and Providers

- Feeding recommendations are the same for all infants, regardless of HIV exposure or HIV infection status.
  - Give only breast milk up to age 6 months.
  - Gradually start complementing breastfeeding with suitable hygienically prepared foods from age 6 months (such as Likuni Phala, fruits, vegetables, beans, ground nuts and soya).
  - Aim to stop breastfeeding around age 22 months, so that the final HIV test can be done at age 24 months (6 weeks after breastfeeding has stopped).
  - Stop breastfeeding gradually over a period of 1 month (no *rapid cessation*).
- 
- Replacement feeding (formula) is **NOT** recommended unless women are unable to breast feed.
  - Monitor weight, height and MUAC according to schedule using standard MOH charts and intervene if no adequate weight-gain.
  - Give only medicines prescribed by a health professional.
  - Start breastfeeding immediately after birth. Explain and observe optimal breastfeeding:
    - Empty both breasts properly to avoid breast engorgement.
    - Ensure proper attachment and positioning to minimize nipple cracks and fissures.
    - Watch out for signs of breast infection (pain, swelling, heat, redness)
      - Don't feed baby from infected breast. Express infected breast to avoid engorgement. Discard expressed milk – do not feed to baby.
      - Go to health facility for treatment. See **section 12.2.8, page 288** in the Malawi Standard Treatment Guidelines (**MSTG 2015**, 5<sup>th</sup> Edition)

## 15.3 Infant NVP prophylaxis



### Key Facts for Patients and Providers

- NVP syrup is given to all babies born to HIV infected mothers.
  - NVP syrup shields the baby from HIV infection during the most risky time.
  - Give NVP syrup to the baby 24-hourly for 6 weeks.
  - All babies should take NVP syrup for the same duration regardless of the mother's ARV regimen and regardless if the mother was taking ARVs at all.
- Store NVP syrup bottles and syringe: dark, cool, clean and dry and out of children's reach.
- Use an old syrup bottle filled with water to show how to draw 1.5ml of syrup in the syringe.
- Hand out one example syringe where the 1.5ml line has been marked with a pen.
- Squirt the syrup in the back of the infant's mouth between the cheek and the gum to ensure it gets swallowed (use cup to demo).
- Rinse the dosing syringe carefully with clean water after every use and let dry.
- Bring back to the health facility at the 6 week vaccination visit all NVP bottles (whether used or unused). The nurse will check if the right amount was used.

### 15.3.1 Prescription and dispensing of NVP prophylaxis

- When to dispense NVP syrup for infant prophylaxis to take home:
  - At ANC (or maternity) as soon as the mother is known to be HIV-infected.
  - Unopened bottles of NVP syrup have a long shelf-life. Therefore never delay dispensing until later in pregnancy. Make sure the expiry date is at least 2 months after the estimated delivery date.
  - Ask at every following visit if the NVP syrup and the syringes are still available. Replace without delay any items that may have been lost or spoilt.
- Dispense 2 x 100ml-bottles of NVP syrup with dosing syringe.

### 15.3.2 Dosing

- The dose of NVP syrup remains the same for the whole 6 week period – do not change the dose according to age or body weight, etc.
- Use the standard dose (1.5ml) If birth weight is unknown (home birth / no scale).

Table 15: Dosing of NVP syrup for infant prophylaxis

Birth weight	NVP syrup (10mg per ml)
2500g or less	1.0 ml 24-hourly
Over 2500g / unknown	1.5 ml 24-hourly

### 15.3.3 Timing and duration

- Start giving NVP syrup as soon as possible after birth. The earlier the start, the more effective.
- NVP syrup can be started anytime between birth and 4 weeks of age if the mother presents late. Starting NVP prophylaxis later is less effective and may cause drug-resistant HIV if the baby is already infected (and needs to start ART).
- Stop giving NVP syrup when the infant is 6 weeks old. The infant will receive less than 6 weeks of prophylaxis if NVP syrup has been started late.

## 16 Pre-exposure prophylaxis (PrEP)



### Key Facts for Patients and Providers

- Other than for HIV exposed infants, the national HIV program does currently **not** support the use of ARVs for pre-exposure prophylaxis.
- Although HIV negative people who are at very high risk of getting infected with HIV can benefit from PrEP, significant implementation challenges have limited the effectiveness in comparable settings.
- Experience from other countries in the region has shown that it is difficult to identify who will be at such high risk that they benefit from PrEP, to keep them in follow-up and to ensure adequate adherence.

## 17 Post exposure prophylaxis (PEP)



### Key Facts for Patients and Providers

- HIV infection can be prevented after a high risk contact with fluids from an HIV infected person.
  - Remove immediately as much as possible of the body fluid.
  - Immediately give a 3-day supply of PEP and start taking it as soon as possible.
  - Assess risk and test for HIV as soon as possible. Continue a 30-day course of ARV prophylaxis (PEP) if exposure is classified as 'risk' and exposed person is HIV negative.
- PEP, if taken correctly, reduces the risk of infection by 80%.
- ARVs taken for PEP are usually well tolerated.
- Keep ARVs for PEP accessible 24/7, e.g. at maternity or other well-advertised locations.
- Offer STI treatment and emergency contraception, for rape victims accessing PEP.
- The risk of getting infected may be high or low, depending on the type of substance and contact. However, PEP should always be started if there is a possible risk of transmission (see classification in **Table 16** on **page 86**).

### Classification of risk

- Use **Table 16** to find out if the exposure is a possible risk for infection.
- Obtaining a new HIV test from the source person can help to reassure that the risk is low, but PEP should still be given if the test result is negative. The source person could be newly infected himself and may be in the window period.

**Table 16: Classification of risk of transmission after exposure to HIV**

	Substance	Type of contact	Source person
<b>Risk</b>	<ul style="list-style-type: none"> <li>• Blood</li> <li>• Semen</li> <li>• Vaginal fluid</li> <li>• Cerebro-spinal fluid</li> <li>• Pleural fluid</li> <li>• Amniotic fluid</li> <li>• Synovial fluid</li> <li>• Ascites fluid</li> </ul>	<ul style="list-style-type: none"> <li>• Skin penetrated with contaminated needle (hollow or non-hollow)</li> <li>• Large amount of substance on mucous membrane</li> <li>• Sexual intercourse no condom</li> <li>• Risk substance on lacerated skin / open wound</li> </ul>	<ul style="list-style-type: none"> <li>• Regardless of known/unknown HIV status</li> </ul>
<b>No Risk</b>	<ul style="list-style-type: none"> <li>• Urine</li> <li>• Stool</li> <li>• Pus</li> <li>• Tears</li> <li>• Saliva</li> <li>• Sputum</li> <li>• Nasal secretions</li> </ul>	<ul style="list-style-type: none"> <li>• Risk substance on intact skin</li> </ul>	

## Immediate measures

- Remove infectious substance.
  - Wash exposed wounds and skin sites thoroughly with soap.
  - Flush mucous membranes with water.
  - Do not use bleach, antiseptics or other caustic substances.

## Eligibility to start PEP (ARV prophylaxis)

- Any exposure classified as risk in the last 72 hours (see **Table 16**).
- Never refuse PEP on moral judgement about the kind of exposure (accident, negligence, rape, 'burst condom').
- New HIV test is mandatory to confirm negative HIV status,
  - BUT: Don't delay starting PEP if HIV testing is not immediately available (no test kits, night, etc.). Do HIV testing as soon as possible.
- PEP is safe in pregnancy and breastfeeding.
- Severe anaemia (<8g/dl) is contraindication for AZT/3TC.
- Severe renal failure is contraindication TDF/3TC.

## How to start PEP

- Start taking PEP as soon as possible after high risk exposure, ideally within 2 hours.
- Starting PEP more than 72 hours after exposure is not effective and should not be done.
  - However, still perform HIV testing at baseline, at 3 and 6 months.
- Explain dosage and importance of adherence.
- Mild side effects (nausea, etc.) are not a reason to stop PEP.
- Advise to return immediately if serious side effects are suspected.
- Advise all exposed adults to practice safe sex until confirmed HIV negative at 3 months.
  - Give 30 condoms and re-supply as requested.
- Do not stop breastfeeding.
- Write case details in PEP register (improvised).

Table 17: Post exposure prophylaxis regimens and dosage (number of tabs taken)

Weight	Standard				Alternative			
	AZT 60mg / 3TC 30mg		AZT 300mg / 3TC 150mg		TDF 300mg / 3TC 300mg		ABC 60mg / 3TC 30mg	
3.0 – 5.9 kg	1	1					1	1
6 – 9.9 kg	1½	1½					1½	1½
10 – 13.9 kg	2	2					2	2
14 – 19.9 kg	2½	2½					2½	2½
20 – 24.9 kg	3	3					3	3
25 – 34.9 kg			1	1				1 1
≥ 35.0 kg					0	1		1 1

### PEP follow-up

- At 30 days: (after completing ARV prophylaxis)
  - Assess adherence
  - Give 60 condoms
- At 3 months and 6 months: repeat HIV testing

### Additional prevention measures after rape / sexual exposure

- Give emergency contraception (EC) within 72 hours if needed (see **Table 18**)
  - Repeat dose if vomiting occurs within 1 hour of taking EC.
  - Explain that next menstrual period should occur before or around the expected time.
- Consider giving presumptive treatment for STIs using **Table 19**

Table 18: Regimens and dose for emergency contraception

Contraceptive drug	Immediately	After 12 hours
Postinor 2 (750µg levonorgestrel)	2 tablets	
OR		
Lo-Feminal or Microgynon	4 tablets	4 tablets

**Table 19: Dosing of standard presumptive STI treatment after sexual exposure**

STI drug	Child <15 years	Adult
Benzathine pen. vials	50,000 IU/kg IM stat (max 2.4 million IU)	2.4 Mega Units IM stat
Gentamicin vials	7.5 mg/kg IM stat (max 240mg)	240mg IM stat
Erythromycin tabs	12.5 mg/kg 6-hourly for 14 days (max 500 mg per dose)	500mg 6-hourly for 7 days
Metronidazole tabs	5 mg/kg 8-hourly for 7 days (max 2 g per day)	2g stat
Nystatin pessaries	N/A	100,000 units 12 hourly for 7 days

## 18 Monitoring and evaluation



### Key Facts for Patients and Providers

- The HIV program relies heavily on accurate and timely data for planning, reporting to donors and for drug procurement and distribution.
- Data analysis and reporting is done from patient cards and clinic registers at most facilities, but electronic systems for monitoring are used at sites with many patients.
- Reporting is done monthly for ANC, maternity and exposed child follow-up and quarterly for ART (see Table 20 on page 93)
- Cohort analyses are needed to report outcomes of patients in ANC, exposed child and ART follow-up. Cohort reports look at the current / latest status of all patients enrolled in follow-up and require a review of all patient records to classify primary and secondary outcomes before data can be aggregated for reporting.
- Reports from facilities are to be completed within 5 working days after the end of the reporting period.
- HIV Program reporting will be further integrated into the regular Health Management Information System. Monthly / quarterly facility reports will be entered directly into the District Health Information System at the District Health Offices for national reporting.

### 18.1 Definitions

#### PMTCT site

- A facility is counted as a PMTCT site if they have initiated on ART at least one pregnant or breast feeding woman during the reporting period.
- Depending on the mode of integration of PMTCT/ART interventions into the general health services, ART may be initiated in any of the following service points: ART, ANC, maternity, post-natal or under 5 clinic.

#### ART site

- A facility is counted as an ART site if they had retained at least one patient alive on ART at the end of the reporting period.

#### ART status at registration

- Refers to the patient's status at the time of first registration at this ART clinic – this status will never change as long as the patient remains at this clinic.
- **First time initiation:** Never taken ART (triple ARV combination treatment) in the past. Having taken ARVs for prophylaxis (PEP, single dose nevirapine, AZT combination prophylaxis for PMTCT) does NOT count as having taken ART and is ignored for the *ART status at registration*.

- **Re-initiation:** Received ART (triple ARV combination for treatment) from another ART site in the past but has NOT been taking it for 2 weeks or more as of the day of registering at this clinic. Patients who have interrupted for 2 weeks or more need to take a starter pack for re-initiation (if started on a regimen containing NVP).
- **Transfer in:** Received ART from another ART site in the past and is currently taking ART or has interrupted for less than 2 weeks. Count as *Transfer In* regardless if the patient brings his old patient card or not ('official' or 'unofficial' transfer).

### Defaulted / Lost to follow-up

- Patients are counted as 'defaulted' in the cohort report if they have not returned to the clinic and are not known to have transferred out, stopped or died.
- The following times apply in the different clinics:
  - HCC (HIV exposed children): 2 months after the *Next Appointment Date* given at the last visit.
  - ART: 2 months after the patient is expected to have run out of ARVs.
- Patients may revert to 'alive on ART' when the next cohort analysis is done if they return to the clinic and continue ART.

### ART stop

- Patients are counted as 'stopped' if they are last known to be alive and have stopped taking ART. Stop is used regardless:
  - Of the reason the patient has stopped (clinician's or patient's own decision).
  - If the ART interruption is intended to be permanent or temporary.
  - Of the duration of the ART interruption at the time of doing the cohort analysis.
- Patients may revert to 'alive on ART' at the next cohort analysis if they re-start ART.

### Died

- Patients are counted as 'died' if there is a reliable report about the patient's death. 'Died' is used regardless:
  - Of the cause of death (HIV- or non-HIV related disease, accident, suicide or homicide).
  - If the patient was on ART or not at the time of death.

### ART re-start

- Interrupted ART for more than 2 months while registered at the respective ART site. Update the number of re-starts in the ART clinic register whenever the patient re-started ART after defaulting or stopping for more than 2 months (i.e. returns after 'defaulting'). Patients who have interrupted for 2 weeks or more need to take a starter pack for re-initiation (if started on a regimen containing NVP).

### ART adherence level

- Reporting of adherence levels is based on a classification of the number of doses missed at the last visit before the end of the quarter evaluated.

- The translation of the number of doses missed into adherence % depends on the number of days since the last visit. In practice, it is too complicated to consider varying intervals when analysing cohort adherence. Therefore, 2 monthly visits are assumed for all when classifying adherence for reporting.
- Patient who are supposed to take 1 tablet per day (e.g. Regimen 5A) and who have missed more than 3 tablets are classified as 'less than 95% adherent'.
- Patients who are supposed to take 2 tablets per day (e.g. Regimen 1A) and who have missed more than 6 doses are classified as 'less than 95% adherent'.

**Table 20: Overview of M&E systems for integrated HIV program reporting**

Service	M&E tools		Report cycle	Report elements			
	Patient card	Register		New registrations	Cohort outcomes		
					Definition of cohort	Primary outcomes	Secondary outcomes
ANC	–	ANC Clinic Register	Monthly	New first visits	•Registration group (6 months after first ANC visit)	–	(Final status at end of ANC) •HIV test status •On ART
Maternity	–	Maternity Register	Monthly	New deliveries	–	–	–
ART	ART Patient Card (separate cards for paediatric and adult formulations)	ART Clinic Register	Quarterly	Patients newly registered at ART clinics	•Cumulative (all ever registered) •Registration group (survival analysis)	•Alive on ART •Died •Defaulted •Stopped ART •Transferred out	•ART regimen / formulation •Adherence level •Side effects •TB status •On CPT •Using FP
Exposed child FUP	HIV Care Patient Card, Exposed Child Under 24 Months	HIV Care Clinic Register	Monthly	Patients newly registered at HCC	•Birth cohort: children who (would) have turned 2, 12 and 24 months of age	•Alive in exp. child FUP •Discharged uninfected •Started ART •Defaulted •Transferred out •Died	•Age when received DNA-PCR result •Latest HIV status

## 18.2 Reporting of registration data

- For all new patients registered, baseline data (such as age at registration, sex, pregnancy status, clinical stage, etc.) are recorded on patient treatment cards and copied into the clinic register.
- These details do not change over time and tallying of these data needs to be done only once when reporting on new patients registered during the reporting month or quarter.
- *Page summaries* in the clinic registers are filled as soon as each page is full. Count the number of circled values for each column on the page.
- **Monthly or quarterly registration reports** are obtained by adding the page summaries from each page in the respective reporting month or quarter.
- **Cumulative registration reports** are obtained by adding the data from the new monthly or quarterly registration report to the data from the previous cumulative registration report.
- Data elements in most sections should add up to the respective total number of patients registered.
  - Males, non-pregnant females and pregnant females must add up to the total number registered.
  - Age groups must add up to the total number registered.
  - ART status (first time initiations, re-initiations, and transfer ins) must add up to the total number registered.
- Some registration data (such as the number of patients with KS at the time of ART initiation) are counted separately and are not part of a section. These data elements are not expected to add up to the total number registered.

## 18.3 Reporting of cohort outcomes

- *Cohort analyses* are needed to measure outcomes of patients in follow-up.
- In principle, the outcome status of any patient ever registered can change at any time. Therefore, the records of all patients ever registered have to be reviewed each time a cumulative cohort outcome analysis is done. Current outcome data cannot be obtained by addition from the previous quarterly outcome data.
- Patient outcomes are considered as of the last day of the reporting period. Any events (e.g. death) that happened after that day are ignored in the respective cohort analysis, but will be counted in the next report.

### Primary follow-up outcome

- The primary outcome shows if a patient has been retained alive in care or if he has dropped out and why.
- The primary outcome categories must add up to the total patients registered in the cohort.
- **Table 20** lists the primary follow-up outcomes used for the different reports.
- For ART only, deaths are further classified according the time after ART initiation. The categories used are: death within 1st, 2nd, 3rd month after ART initiation or after 3rd month of ART initiation.

## Secondary outcome

- Secondary outcomes are the latest treatment details among the patients retained alive in care.
- Secondary outcomes are counted directly from the cards of the patients retained alive in care, usually by looking at the last visit before the end of the month or quarter evaluated. This visit might be several months before the end of the quarter, for example if the patient is on long ARV dispensing intervals (as long as the patient is still classified as 'retained alive in care' at the end of the quarter evaluated).
- Each set of secondary outcome categories must add up to the total number of patients retained alive in care.
- **Table 20** shows the secondary outcomes used for the different reports

## Definition of cohorts for different program reports

- 3 slightly different methods are used to define cohorts for outcome analyses:
- **Cumulative cohort** (ART): Follow-up status of all patients ever registered at the respective clinic. The number of patients with adverse follow-up outcomes (death, default, etc.) inevitably increases over time. The number of patients retained in care is calculated by subtracting all patients with adverse follow-up outcomes from the total patient ever registered.
- **Registration group cohort** 'Survival analysis' in ART: Follow-up status of patients registered during the quarters that ended 12, 24, 36, 48 and 60 months ago (ART). *ANC cohort outcomes*: final status as of the last ANC visit for the women who started ANC 6 months ago. This method standardises follow-up times and makes outcome data comparable between sites and over time.
- **Birth cohort** (HIV exposed child follow-up): Follow-up status of children who (would) have turned 2, 12 and 24 months old. Patient cards are filed in batches by month and year of birth (birth cohorts) and only the cards of children born 2, 12 and 24 months ago are pulled out for reporting. Outcomes are counted separately for the 2-, 12- and 24-month birth cohort. Reporting is done monthly and a different birth cohort is covered in each reporting month. This method standardises ages and is used for children enrolled in HIV exposed child follow-up.

## 18.4 Record keeping and filing

### Confidentiality of patient records

- All patient cards and clinic registers are property of the MOH and may only be kept at the respective facility or at the National Archives.
- Patient cards and clinic registers must be kept in a locked room and are only to be accessed by clinic staff responsible of providing the respective service and by the national supervision team. Patients and named guardians have access to their own patient card.

### Use of clinic registers (ANC, Maternity, HCC, ART)

- Keep patient registration for each different service centralized in each facility: Use only one set of registers in each facility.
- Each patient has only one row<sup>14</sup> in each register: Continue using the same row for returning transfers and re-starts after default or stop.

<sup>14</sup> In the ANC register, each woman has one separate section with rows for each subsequent visit.

- Turn to a new page when starting to register patients in a new quarter. Leave any unused rows at the bottom of the previous page empty. This is to separate the quarters when adding page totals.
- Assign continuous registration numbers (by sequence of registration). Take care not to duplicate registration numbers.
  - Continue assigning cumulative registration numbers in the HCC- and ART-Register. These number series are never re-started.
  - Re-start assigning registration numbers annually for the ANC- and Maternity Register. Re-start with number 1 on the 1<sup>st</sup> of July.

### Use of patient cards

- Each patient has only one patient card at any one time (Exposed child, ART). Attach another patient card once the old card is full.
- Patient cards are filed in polythene sleeves in lever arch files, up to 100 cards per arch file.
- Separate filing systems are used for the different types of patient cards:

#### Exposed Child under 24 Months cards

- File in batches by year and month of birth.
- Within each birth month, sort in ascending order by HCC registration number.
- Do not remove the cards of children who have started ART, died, defaulted or transferred out from this filing system.
- Files with birth cohorts who (would) have now reached at least age 3 years can be removed from the clinic for archiving.

#### ART Patient cards, paediatric and adult ARV formulations

- File ART Patient Cards in ascending order by ART registration number.
- Prepare separate filing systems for **ACTIVE** (retained in ART) and **INACTIVE** patients (stopped ART, transferred out, defaulted, died).
- One arch file can hold approximately 100 cards.
  - Label the **ACTIVE** files with ART numbers 1-100, 101-200, 201-200, etc.
  - Label the **INACTIVE** files with ART numbers 1-200, 201-400, 401-600, etc.
- Each time the quarterly cohort analysis is done, update in the ART register the outcome for patients who have dropped out of ART (stopped ART, transferred out, defaulted or died). Straight after this, move these cards of from the **ACTIVE** to the **INACTIVE** filing system.
- Do not separate paediatric and adult ARV formulation cards into different files.

## 18.5 Ensuring adequate data quality

- Use only the standard national reporting forms.
- The clinic's own reports are checked by the supervision team each quarter from primary records.
- Copies of the checked reports are kept at the clinic.

## 19 Supply Management



### Key Facts for Patients and Providers

- The HIV program requires an uninterrupted supply of huge amounts of very expensive drugs and lab supplies. Between 2016 and 2017, Malawi will spend over USD 300 million (MKW 166 billion) for HIV commodities.
- Commodity stock-outs lead to an interruption of life-saving health services. ARV stock-outs are especially serious because patients who interrupt treatment can develop drug-resistant HIV which can be transmitted to others.
- A physical buffer-stock of HIV program commodities is maintained at the central warehouse to ensure uninterrupted supply to facilities. The buffer stock is maintained to cover to **6 months** consumption.
- **Responsibilities:**
  - All health workers: support supply management by filling the standard MOH forms, patient cards, registers and reporting forms.
  - Officer in-charge of pharmacy: manage and account for all commodities received.
  - District Health Management Teams: coordinate and supervise.
- A dedicated HIV Program Logistics Team (**HIV Logistics**) working under MOH Depts. for Health Technical Support Services and for HIV and AIDS actively coordinates procurement, supply planning and distribution of medicines and lab supplies for the HIV and STI Programs.
- Contact HIV Logistics by email ([hivdeptlogistics@gmail.com](mailto:hivdeptlogistics@gmail.com)) or call **toll-free** on working days 7:30 – 16:30:
  - **5 91 91** (from Airtel phone)
  - **68 82** (from TNM phone)
- Ask HIV Logistics for help and get an **authorization code before** any of the following transactions with ARVs and HIV test kits:
  - Getting additional supplies from warehouse.
  - Moving stocks from / to another facility.
  - Disposing expired / spoiled stocks.
- **Notify** HIV Logistics about (even if suspected):
  - Damaged or inappropriate stocks received.
  - Serious (suspected) side effects.

## 19.1 HIV commodity supply cycle

- Table 21 shows the different commodity groups currently managed by the HIV Program.

**Table 21: Drugs and testing supplies managed by the HIV Program**

Commodity group	Examples	Supply*
ARVs	(All ARVs, incl. PEP and infant prophylaxis)	E
OI	Cotrimoxazole for CPT	E
	Isoniazid + pyridoxine for IPT	E
	Cotrimoxazole, other antibiotics, fluconazole, chemotherapy	S
STI	Standard / alternative antibiotics, acyclovir, clotrimazole	S
PIFP	Condoms, Depo-Provera	S
Analgesic	Morphine, codeine	S
DBS kits	for EID and VL samples	E
Tests	HIV and syphilis rapid test kits	E

Supply\*: **E** = item managed exclusively through HIV Program. **S** = items supplemented by HIV Program in addition to essential medicine supplies.

- HIV commodities are delivered in the every 2 months from a central warehouse (Lilongwe) directly to all facilities.
- Distribution lists for all facilities are calculated based on the patient and stock reports collected during quarterly HIV Program supervision and reported through the Logistics Management Information System.
- Actively support the **2-monthly supply cycle** and the **ongoing management** following the 11 steps in **Figure 7**.


### 1. Prepare stock / patient report

- Confirm each commodity is sorted by expiry date.
- Do physical count of stock on hand (SOH). Exclude any units that may have already expired.
- Ensure all available stock is counted, including in bulk store, at the clinic / HIV testing rooms, etc.

### 2. Verify data

- Ensure all storage areas and patient records / registers are accessible on the day of supervision.
- The HIV Program supervision team will work with facility staff to verify:
  - Stock reports by doing a physical count.
  - Patient report by reviewing patient cards and registers.
- Check that the stock report filled during supervision is complete and accurate. The supervision team and the In-Charge of pharmacy are responsible for confirming this by signing the form.

Figure 7: Flowchart for HIV commodity supply management



### 3. Review draft distribution list

- **2-monthly consignments** are calculated by *HIV Logistics* from patient numbers and stock reports collected at the last supervision visit.
  - All ARVs and HIV test kits included should be about **2 MOS** (months of stock, see below).
  - Consignments are scheduled to arrive in every 2 months.
  - Facilities should have about **2 MOS** remaining when the new consignment arrives (site-level buffer). This should bring the total to about **4 MOS**.
- HIV Logistics will circulate the *draft distribution list* to anyone registered with their email address. To register, send an email request to [hivdeptlogistics@gmail.com](mailto:hivdeptlogistics@gmail.com). Anyone can also subscribe for an automatic notification by email and/or SMS whenever a distribution list is posted for review on the HIV Dept. website ([www.hiv.health.gov.mw](http://www.hiv.health.gov.mw)).
- Review and confirm that the items and quantities are correct and adequate for you. Submit any suggested changes (by email, SMS or phone) before the deadline shown on the draft list.

### 4. Receive consignment

- Inspect the entire consignment in the presence of a witness designated by DHMT/ facility In-Charge:
  - Physically count all re-packed / loose units. Originally sealed boxes do not need to be opened for counting of units. Add up total units received for each item.
  - Check expiry date for all items.
  - Write physical count for each item into the respective box on the *delivery note*. Write 0 (zero) for any items not received – don't leave any boxes empty.
- Sign, date and stamp the *delivery note* to confirm receipt of the items as indicated.
- The person signing on the delivery note is accountable for all items s/he has signed for. The In-Charge of pharmacy / facility will be held responsible for any discrepancies noted later.

### 5. Store

- Immediately move all items received to a secure storage area (clean, dry, cool and off the floor).
- Enter quantity and date of receipts on *stock cards* without delay.
- Arrange items by expiry date to make it easy to follow the *First Expiry -First Out* principle (*FEFO*).

### 6. Issue to clinic

- Fill *Requisition and Issue Vouchers* for all commodity requests from the clinic.
- Always follow the *FEFO* principle.
- Immediately update *stock card* when moving items out of the pharmacy.
- Limit the amount of stock stored at the clinic to 1 week consumption.

### 7. Dispense / use

- Ensure that the patient has fully understood:

- How and when to take their drugs.
- Possible side-effects; which side-effects require coming to the health facility.
- Account for all HIV commodities dispensed. Specify type and quantity:
  - On *patient master cards* (ART, Exposed child)
  - *Dispensing registers* for special drugs (Diflucan)
  - *Daily Activity Registers (DAR)* for HIV test kits.
- The *DAR* is used for tracking use of HIV test kits.
  - Keep a separate register at all places where HIV testing is done.
  - Use separate pages for the different types of tests (Determine, Uni-Gold).
  - Test kits *used for clients* must match entries in the HIV testing Register.
  - The DAR includes sets of 3 carbonated sheets: keep white sheet at facility; send blue sheet to DHO; retain pink/yellow sheet for collection by HIV Logistics (MOH).
  - Fill monthly summary on HIV testing report by adding numbers from all DAR used at the facility.

## 8. Monitor stocks / consumption

- Do a **physical stock count** for all items (in store and at the clinic) and update stock cards:
  - On the last working day of each month.
  - When handing over pharmacy management to another staff member.
  - Whenever discrepancies are noted.
- **Calculate** average monthly consumption (**AMC**) and months of stock (**MOS**) for all ARVs and HIV test kits after doing the monthly physical count:

$$\text{AMC} = \frac{\text{units used in last 3 months}}{3} \qquad \text{MOS} = \frac{\text{stock on hand}}{\text{AMC}}$$

- Be alert: commodity shortages can be anticipated before they happen:
  - Large number of transfers in.
  - Patients moving to 2<sup>nd</sup> line or alternative regimens.
  - Rapid growth through new initiations.
- As soon as commodity **shortage** is **suspected or noticed**:
  - Contact *HIV Logistics* for additional supply (see below).
  - Inform all relevant staff members.
  - Prioritize use (e.g. HIV test kits for sick patients needing to start ART, women at ANC and maternity, etc.).
  - Shorten supply interval (e.g. give ARVs for 1 month instead of 3).
- Commodity **excess**: more than **4 MOS**, especially if units will expire before they can be used:
  - Contact *HIV Logistics* for to request stock relocation (see below).

## 9. Request adjustment

- Call *HIV Logistics* as soon as possible if **shortage**, **excess** or **expiry** is noted.
- Before calling, prepare the following information:
  - Number of tins / bottles / tests remaining.
  - Expiry date
  - Number of patients on this regimen / approximate AMC.
  - When additional stocks are needed / to be sent to other site.
  - If own transport can be organized.
- *HIV Logistics* will:
  - Review the information and find out the reason for the problem.
  - Coordinate: **extra allocation from the warehouse**, **relocation of stocks between sites**, or register **disposal of expired commodities**.
  - Send a unique **Authorization Code** for each item by SMS or phone.
- Confirm receipt of authorization codes by sending 'OK' by SMS or by calling *HIV Logistics*.
- Fill a **Registration Form for Relocation or Disposal of HIV Commodities** for each adjustment.
  - Write the authorization code for each item on the form.
  - Keep Registration Forms in the pharmacy to account for all commodity transactions.
- Never relocate or dispose HIV commodities without *authorization code*. In exceptional circumstances (threatening stock-out and no phone coverage / no answer), stocks may be relocated and notification and *authorization codes* must be obtained at the earliest opportunity.

## 10. Collect / receive / release stock (from adjustment)

- When collecting extra consignments from the warehouse:
  - Ask for the size of the consignment and make sure it can be safely transported (security, sun/rain protection, etc.). Partial collection will not be allowed.
  - Make specific appointment and get directions from *HIV Logistics*.
  - Bring ID (passport, driving license, etc.) and official facility stamp.
  - Inspect the whole consignment. The collecting officer and a witness must fill, sign and stamp the delivery note as usual.
  - There is no need to fill a *Registration Form for Relocation* for extra allocations from the warehouse.
- Relocating stocks between facilities:
  - Fill a *Registration Form for Relocation* and write the *authorization code* for each item.
  - Keep the white copy of the form at the facility releasing the stock. This is mandatory to account for commodities given away to another site.
  - Give the pink copy to the facility receiving the relocated commodities.

## 11. Manage disposal

- Separate expired commodities from usable stock as soon as possible.
- Notify HIV Logistics, get **Authorization Code** and fill **Registration Form for Disposal**
- Contact the District Pharmacist and arrange for transfer of expired items for controlled destruction.

# Index

3TC.....	43	table .....	48
ABC/3TC		third line .....	42
dosing .....	50	ART registration status definitions .....	90
abdominal pain		ART re-start (M&E definition) .....	91
differential diagnosis .....	32	ART site definition .....	90
differential diagnosis and management .....	78	ART stop (M&E definition) .....	91
active patient file .....	96	artificial rupture of membranes .....	81
adherence		ARV classification .....	43
appointment .....	66	ARV formulations	
calculation .....	53	adult and paediatric formulations .....	44
classification for reporting .....	91	dosing table .....	50
counseling .....	68	ascertainment of HIV exposure in children .....	15
dose .....	68	flowchart .....	16
age cohort .....	95	ascertainment of HIV status .....	15
amitryptilline .....	79	<b>ATV/r</b>	
anaemia		Atazanavir / ritonavir .....	III, 31, 43, 49, 51, 64, 65
ART regimen substitution .....	48	AZT/3TC	
starting ART .....	51	dosing .....	50
ANC		AZT/3TC/NVP	
standard HIV services delivered in .....	11	dosing .....	50
ANC register .....	95	BCG vaccination .....	82
appointment scheduling .....	53	birth cohort .....	95
archiving of patient records .....	95	bleomycin .....	21
ARM .....	81	blisters	
ART		differential diagnosis .....	33
adherence classification .....	91	BMI	
alternative regimens .....	48	classification .....	28
'emergency dispensing' .....	53	definition .....	27
ART clinic		<b>body pains</b>	
standard HIV services delivered in .....	11	differential diagnosis and management .....	76
ART clinic register .....	95	breast enlargement .....	See gynaeomastia
ART contraindications .....	45	breast infection .....	83
ART counselling		breastfeeding	
group .....	62	counseling .....	83
individual .....	62	buffalo hump .....	30
ART dispensing		bumpy itch .....	See pruritic papular eruptions
calculation of quantities .....	53	cancer .....	78
maximum supply .....	54	CD4 monitoring	
patients on 2nd line regimens .....	54	sample collection .....	34
safety buffer .....	54	schedule .....	34
supply and adherence calculation (table) .....	55	cervical cancer	
ART duration, monitoring of .....	71	diagnosis and management .....	21
ART failure		clinical monitoring checklist .....	29
confirmed .....	73	cohort analysis .....	93
suspected .....	71	definition of cohorts for reporting .....	95
ART initiation .....	56	registration groups .....	95
counseling .....	61	condoms	
labour .....	82	routine dispensing .....	37
preparing the patient .....	61	confidentiality of patient records .....	95
ART patient card .....	57	confirming HIV infection .....	57
adult ARV formulation .....	96	testing protocol .....	58
paediatric ARV formulation .....	96	<b>confusion</b> .....	79
ART prescriber level .....	44	cotrimoxazole preventive therapy .....	37
ART regimen		<b>cough</b>	
selecting regimen, formulation and dosage .....	43	differential diagnosis .....	31
selection for continuation .....	67	CPT .....	37
ART regimens .....	42	contraindications .....	38
first line .....	42	duration .....	38
second line .....	42	eligibility .....	38

cryptococcal meningitis		
management .....	23	
cumulative reporting		
outcome data .....	95	
registration data .....	94	
d4T/3TC/NVP		
dosing .....	50	
DBS		
Dried blood spot .....	III, 34, 58, 72	
death (M&E definition) .....	91	
defaulter definition .....	91	
delivery		
management .....	81	
diarrhoea		
differential diagnosis .....	33	
differential diagnosis and management .....	77	
diarrhoea, chronic		
management .....	20	
died (M&E definition) .....	91	
dizziness .....	29, 30, 48, 61, 67	
doses missed		
classification for reporting .....	91	
doses missed (ART) .....	66	
drowsiness .....	79	
efavirenz		
skin rash .....	48	
teratogenicity .....	45	
EFV		
dosing .....	50	
emergency ARV dispensing register .....	53	
emergency contraception .....	88	
emergency dispensing (ART) .....	53	
exposed child follow-up		
enrolment .....	82	
Exposed child follow-up		
standard HIV services delivered in .....	11	
exposed child under 24 months patient card .....	96	
failure to thrive		
differential diagnosis .....	30	
family planning .....	36	
Family planning clinic		
standard HIV services delivered in .....	11	
fattening		
diagnosis and management .....	76	
FDC .....	43	
fever		
differential diagnosis .....	32	
differential diagnosis and management .....	76	
filing systems (M&E) .....	95	
first time initiation (ART registration status) .....	90	
fixed dose combination .....	43	
glands, swollen		
differential diagnosis .....	30	
growth curve .....	27	
gynaecomastia .....	48	
hanging tablets .....	54	
HCC register .....	95	
hepatitis .....	78	
ART regimen substitution .....	48	
starting ART .....	51	
herpes simplex .....	31	
herpes zoster		
management .....	21	
HIV dementia .....	30	
HIV exposure		
risk classification .....	86	
HIV-related diseases		
clinical management .....	19	
monitoring .....	66	
immune reconstitution inflammatory syndrome .....	80	
implementation plan .....	6	
inactive patient file .....	96	
infant and child feeding .....	11, 13, 83	
In-patient Dept.		
standard HIV services delivered in .....	11	
insecticide treated bed nets .....	11, 13, 41	
integration		
clinical HIV services .....	10	
standard package of services .....	11	
interaction		
ART and TB treatment .....	64	
IPT .....	39	
contraindications .....	40	
dosage and duration .....	40	
eligibility .....	40	
ruling out of active TB .....	40	
IRIS .....	80	
differential diagnosis .....	80	
management .....	80	
isoniazid preventive therapy .....	11, 13, 39	
ITN .....	See insecticide treated bed nets, See insecticide treated bed nets, See insecticide treated bed nets	
jaundice		
differential diagnosis .....	31	
starting ART .....	51	
Kaposi sarcoma		
management .....	21	
kidney failure .....	See renal failure	
lab investigations .....	63	
labour		
management .....	81	
starting ART .....	51	
lactic acidosis .....	77	
lamivudine .....	43	
leg pain		
differential diagnosis .....	33	
differential diagnosis and management .....	79	
liquid suspension (ART) .....	45	
loss to follow-up (ART) .....	75	
loss to follow-up definition .....	91	
LPV/r		
dosing .....	50	
lymphadenopathy		
differential diagnosis .....	30	
M&E .....	90	
malaria .....	78	
master cards .....	96	
Maternity		
standard HIV services delivered in .....	11	
maternity register .....	95	
monitoring and evaluation .....	90	
mouth sores		
differential diagnosis .....	31	
MUAC		
mid upper arm circumference in pregnant women .....	27	
nausea		
differential diagnosis .....	30	
differential diagnosis and management .....	77	
neck stiffness		
differential diagnosis .....	30	

needle stick injury.....	86	rape.....	88
nephropathy		rash	
differential diagnosis .....	78	differential diagnosis.....	33
nevirapine prophylaxis .....	84	registration data (reporting) .....	94
dispensing.....	84	registration numbers .....	96
dosing .....	84	re-initiation (ART registration status).....	91
timing and duration .....	85	renal failure	
newborn care .....	82	ART regimen substitution.....	48
night sweats		starting ART .....	51
differential diagnosis .....	32	Renal failure .....	38, 48, 49
nightmares.....	79	reporting	
<b>NNRTI</b> .....	43	outcome data .....	94
<b>NRTI</b> .....	43	registration data.....	94
nutrition status		reporting cycle .....	93
BMI .....	27	re-start ART (M&E definition) .....	91
monitoring.....	27	reverse transcriptase inhibitor.....	43
nutritional monitoring .....	66	seborrhoeic dermatitis	
oesophageal candidiasis		management .....	22
management .....	20	secondary outcome.....	93, 95
OPD		sepsis	
HIV services delivered in .....	11	management .....	23
OPV.....	82	shingles	
oral candidiasis		management .....	21
management .....	19	shortness of breath	
oral hairy leukoplakia .....	31	differential diagnosis .....	32
oral polio vaccination .....	82	skin rash	
outcome		ART regimen substitution.....	48
ART follow-up .....	75	differential diagnosis and management.....	77
outcome data (reporting) .....	94	slimming.....	30
page summary .....	96	differential diagnosis and management.....	76
page summary (reporting).....	94	standard monitoring of HIV patients.....	27
patient cards.....	96	starter pack	
PCP		rationale .....	44
management .....	22	schedule of next appointment .....	54
PEP.....	86	when to give and when not to give .....	44
peripheral neuropathy		Stevens-Johnson-Syndrome.....	33, 45
ART regimen substitution .....	48	STI treatment, presumptive .....	89
differential diagnosis .....	33	stopping ART	
differential diagnosis and management .....	79	indications .....	66
PI 43		substitution (ART regimen)	
PIFP .....	36	strategy .....	45
pill count (ART) .....	66	survival analysis.....	95
pleural effusion		swelling (face)	
differential diagnosis .....	32	differential diagnosis and management.....	78
PMTCT site definition .....	90	tail (stopping ART)	
pneumocystis carinii (jiroveci) pneumonia (PCP)		rationale .....	44
management .....	22	when not to give .....	44
pneumonia		TB	
management .....	23	management .....	20
poor adherence .....	68	TB clinic	
post exposure prophylaxis.....	86	standard HIV services delivered in .....	11
eligibility for ARV prophylaxis .....	87	TB screening, routine .....	67
immediate measures .....	87	TB treatment	
postnatal care.....	82	combining with ART .....	64
Postnatal clinic		TB, active	
standard HIV services delivered in .....	11	starting ART .....	51
pregnancy, first trimester		TDF/3TC	
starting ART .....	51	dosing.....	50
presumptive STI treatment.....	89	TDF/3TC/EFV	
preventive services for HIV patients.....	36	dosing.....	50
primary outcome .....	94	tenofovir	
primary outcomes .....	93	bone growth.....	44
protease inhibitors .....	43	tinea corporis / cruris / pedis	
provider initiated family planning .....	36	management .....	22
pruritic papular eruptions			
management .....	22		

toxicity (ART, severe) .....	45	targeted .....	73
transfer in (ART registration status) .....	91	vomiting	
umbilical cord cutting.....	81	differential diagnosis .....	30, 32
Under 5 clinic		differential diagnosis and management .....	77
standard HIV services delivered in .....	11	weakness	
vacuum extraction .....	81	differential diagnosis and management .....	76
veins, protruding .....	30	weight loss	
Vincristine .....	21, 33	differential diagnosis .....	30
viral load		yellow eyes	
monitoring.....	72	differential diagnosis .....	31
monitoring schedule .....	72	differential diagnosis and management .....	78
routine monitoring .....	72		