

Example policy 3

Approved PrEP methods for differentiated PrEP service delivery

The following PrEP methods have been approved, and this guidance provides the differentiated PrEP service delivery model options across these methods.

(insert country approved PrEP methods)

Service delivery constraints and limitations are indicated below by approved method.

a. Oral PrEP (daily and/or event-driven)

- Initiation prescribing and rescripting by *(insert authorised service providers)*
- No refrigeration requirements. Stored safely at room temperature.

b. Dapivirine vaginal ring

- Initiation prescribing and rescripting by *(insert authorised service providers)*
- No refrigeration requirements. Stored safely at room temperature.
- Dapivirine vaginal rings are inserted by the person using PrEP and do not require an in-person consultation for monthly ring insertion.

c. Long-acting injectable cabotegravir

- Initiation prescribing and rescripting by *(insert authorised service providers)*
- No refrigeration requirements. Stored safely at room temperature.
- First two injections four weeks apart. Thereafter, repeat dosing is required every second month.
- Injection administration by *(insert service providers authorised to give intramuscular, gluteal injections)*
- *(insert any oral PrEP required for lead-in or discontinuation as per country PrEP guidelines)*

This policy example is linked to the publication, "IAS - the International AIDS Society. *Key considerations in developing policy guidance for differentiated PrEP service delivery: Country policy development brief.* July 2022." available at <https://www.differentiatedservicedelivery.org/Resources/Resource-Library/DSD-PrEP-policy-2022>

Download a word version of an example differentiated PrEP service delivery policy at <https://www.differentiatedservicedelivery.org/Resources/Resource-Library/PrEP-example-policy-word>